


**Tuesday, April 23** \*\* Stokes' Theorem

1. Let  $S$  be the portion of the cylinder of radius 2 about the  $x$ -axis where  $-1 \leq x \leq 1$ .
  - (a) Draw a picture of  $S$  and compute its area without doing any integrals. Hint: How could you make this cylinder out of paper?
  - (b) Find a parameterization  $\mathbf{r}(u, v)$  of  $S$ .
  - (c) Does the normal vector field associated to your parameterization point into or out of  $S$ ? First, try to determine this without doing any calculations, and then check your answer by evaluating  $\mathbf{r}_u \times \mathbf{r}_v$ .
  - (d) If necessary, change your parameterization so that the normal vector field points *inwards*.
  - (e) Now consider the vector field  $\mathbf{F} = \langle -z, xz, -xy \rangle$ . Compute  $\text{curl} \mathbf{F}$ .
  - (f) Check that  $\text{curl} \mathbf{F}$  is the sum of  $\mathbf{G} = \langle -2x, -1, 0 \rangle$  and  $\mathbf{H} = \langle 0, y, z \rangle$ .
  - (g) Use geometric arguments to determine whether the flux of  $\mathbf{G}$  is positive, zero, or negative. Remember that we have oriented  $S$  so that the normals point *inwards*. Do the same for  $\mathbf{H}$  and  $\text{curl} \mathbf{F}$ .
  - (h) Using your parametrization, directly compute the flux of  $\text{curl} \mathbf{F}$ .
  - (i) Check your answer in (h) using Stokes' Theorem. Note here that  $\partial S$  has two boundary components, and make sure that you orient them correctly.
  - (j) Check your answer in (h) a second time by using what you learned in (g) to compute the flux of  $\mathbf{G}$  and  $\mathbf{H}$ .
  
2. Consider the surface  $S$  shown below, which is oriented using the outward pointing normal.



- (a) Suppose  $\mathbf{F}$  is a vector field on  $\mathbb{R}^3$  which is equal to  $\text{curl} \mathbf{G}$  for some unknown vector field  $\mathbf{G}$ . Suppose the line integral of  $\mathbf{G}$  around the unit circle (oriented counter-clockwise) in the  $xy$ -plane is 25. Determine the flux of  $\mathbf{F}$  through  $S$ .
- (b) Suppose  $\mathbf{H}$  is a vector field on  $\mathbb{R}^3$  which is equal to  $\text{curl} \mathbf{B}$  for some unknown vector field  $\mathbf{B}$ . If  $\mathbf{H}(x, y, 0) = \mathbf{k}$ , find the flux of  $\mathbf{H}$  through the surface  $S$ .

Check your answers with the instructor.