

On calibrated representations and the Plancherel Theorem for affine Hecke algebras

James Parkinson

Received: 19 November 2012 / Accepted: 27 November 2013 / Published online: 24 December 2013
© Springer Science+Business Media New York 2013

Abstract This paper has two main purposes. Firstly, we generalise Ram’s combinatorial construction of calibrated representations of the affine Hecke algebra to the multi-parameter case (including the non-reduced BC_n case). We then derive the Plancherel formulae for all rank 1 and rank 2 affine Hecke algebras, using our calibrated representations to construct all representations involved.

Keywords Affine Hecke algebra · Calibrated representations · Plancherel measure

1 Introduction

In this paper, we extend Ram’s combinatorial construction of calibrated representations of affine Hecke algebras to the multi-parameter case (including the non-reduced case), and we use these representations to derive explicit Plancherel formulae for all rank 1 and rank 2 affine Hecke algebras, following the work of Opdam (see [18, 19]).

Let us discuss the relevance and significance of each of these objectives. Affine Hecke algebras arise in the study of representation theory of groups G of Lie type defined over local fields such as $\mathbb{F}_q((t))$ or \mathbb{Q}_p . If I is an *Iwahori subgroup* of G then complex representations of G with vectors fixed by I can be studied via corresponding representations of the associated affine Hecke algebra $\mathcal{H} = \mathcal{C}_c(I \backslash G / I)$ of continuous compactly supported I bi-invariant complex valued functions on G (see [2, 16]). On the one hand, the representation theory of affine Hecke algebras is well behaved (for example, the irreducible representations of these infinite dimensional algebras are all finite dimensional), while, on the other hand, the representation theory is rather delicate (for instance, see the remarkable geometric classification of the irreducibles given in [12] using the K -theory of the flag variety).

J. Parkinson (✉)
University of Sydney, Sydney, Australia
e-mail: jamesp@maths.usyd.edu.au

Affine Hecke algebras have a basis $\{T_w \mid w \in W\}$ indexed by elements of an *affine Weyl group* W , and depend on parameters q_0, \dots, q_n (one parameter for each Coxeter generator s_0, \dots, s_n of W). The most studied case is the *1-parameter case*, where $q_i = q$ for all i . It is to this case that the geometric classification mentioned above applies. In [23], Ram introduced an explicit combinatorial construction of the class of *calibrated representations* of 1-parameter affine Hecke algebras. These are the modules which have a basis of simultaneous eigenvectors for all the elements of a natural large commutative subalgebra of the Hecke algebra. While not all representations of an affine Hecke algebra are calibrated, the calibrated representations are of particular interest to combinatorialists since they are the generalisation of the classical combinatorial constructions for Weyl groups to (one parameter) Hecke algebras. Our first aim in this paper is to extend the construction of calibrated representations to the multi-parameter case (see Theorem 3.6). We suspect that a full classification of calibrated representations, along the lines of the one parameter case, is possible, although we defer this investigation to later work and instead the focus here is on constructing calibrated representations. In Sect. 3.3, we give some explicit examples of our calibrated representations, and, in Sect. 3.4, we develop the character theory of calibrated representations in preparation for the Plancherel Theorems.

In the second part of this paper, we derive the Plancherel Theorem for rank 1 and 2 affine Hecke algebras. The Plancherel Theorem is the spectral decomposition of the canonical trace functional $\text{Tr} : \mathcal{H} \rightarrow \mathbb{C}$ with $\text{Tr}(T_w) = \delta_{w,1}$ for w in the affine Weyl group W . It is the analogue of the formula

$$\text{Tr}(a) = \sum_{\pi \in \text{Irrep}(\mathcal{H})} m_{\pi} \chi_{\pi}(a)$$

for finite dimensional Hecke algebras, where m_{π} are the *generic degrees* (see [8, Chap. 11]). For affine Hecke algebras the sum becomes an integral over representations of a C^* -algebra completion of \mathcal{H} , and the weights m_{π} become the *Plancherel measure*.

The Plancherel Theorem has been proven in general by Heckman and Opdam [10] and Opdam [19] in a veritable tour-de-force paralleling Harish–Chandra’s work [9] on the Plancherel Theorem for real and p -adic Lie groups (see also Reeder [26]). The Plancherel Theorem has been further developed by Delorme–Opdam, Opdam, Opdam–Solleveld, and Ciubotaru–Kato–Kato (see [4, 6, 20, 21]). Therefore, we should explain the value of our direct calculations in ranks 1 and 2.

Firstly, while the general formulation of the Plancherel Theorem in [19] is essentially complete, there are some constants that are not explicitly computed (they are conjectured in [19, Conjecture 2.27] to be rational numbers). Thus it is desirable to have a complete and direct calculation in ranks 1 and 2 which evaluate all constants involved. (We note that in the case of the affine Hecke algebra of the general linear group over a non-Archimedean local field, the Plancherel Formula is entirely known, see [1, Remark 5.6]).

Secondly, for concrete applications of the Plancherel Theorem (for example, probabilistic calculations like in [22]) one may need explicit constructions of the representations involved in the Plancherel formula. For the non-expert this may be a difficult task to fulfil, and so we believe that the combination of both parts of this paper, with

a very concrete matching up of representations and terms in the Plancherel Theorem, is of value. In particular, the use of calibrated representations makes the Plancherel Theorem accessible at a combinatorial level.

Finally, we hope that the explicit calculations may in some ways serve as an introduction to the general theory, and illustrate the complexity involved in the sophisticated work [4, 19]. The starting point and general philosophy for our derivation of the Plancherel Theorems is similar to that in [19], but since we restrict to the rank 1 and 2 cases the calculations can be carried out by hand. In fact, our calculations form an extension of Matsumoto’s influential rank 1 calculations [16, §2.6], and hence provides a companion piece to [16] (see also Kutzko and Morris [13]).

2 Definitions and setup

2.1 Root systems and Weyl groups

Let R be an irreducible (not necessarily reduced) finite crystallographic root system with simple roots $\alpha_1, \dots, \alpha_n$ in an n -dimensional real vector space V with inner product $\langle \cdot, \cdot \rangle$. Let R^+ be the set of positive roots relative to the simple roots $\alpha_1, \dots, \alpha_n$. Let W_0 be the *Weyl group*; the subgroup of $GL(V)$ generated by the reflections $s_\alpha, \alpha \in R$, where $s_\alpha(\lambda) = \lambda - \langle \lambda, \alpha \rangle \alpha^\vee$ with $\alpha^\vee = 2\alpha / \langle \alpha, \alpha \rangle$. Thus W_0 is a Coxeter group with distinguished generators s_1, \dots, s_n (where $s_i = s_{\alpha_i}$). Let w_0 be the (unique) longest element of W_0 . The *dual root system* is $R^\vee = \{\alpha^\vee \mid \alpha \in R\}$. The *coroot lattice* Q and the *coweight lattice* P are

$$Q = \mathbb{Z}\text{-span of } R^\vee \quad \text{and} \quad P = \mathbb{Z}\omega_1 \oplus \dots \oplus \mathbb{Z}\omega_n,$$

where $\omega_1, \dots, \omega_n$ are the *fundamental coweights* defined by $\langle \alpha_i, \omega_j \rangle = \delta_{ij}$. The cone of *dominant coweights* is $P^+ = \mathbb{Z}_{\geq 0}\omega_1 \oplus \dots \oplus \mathbb{Z}_{\geq 0}\omega_n$. Then $Q \subseteq P$, and W_0 acts on lattices L with $Q \subseteq L \subseteq P$. The *affine Weyl group* associated to R and L is

$$W_L = L \rtimes W_0,$$

where we identify $\lambda \in L$ with the translation $t_\lambda(x) = x + \lambda$. Let φ be the highest root of R , and let $s_0 = t_\varphi s_\varphi$. Let $S = \{s_0, \dots, s_n\}$. Then $W_Q = \langle S \rangle$ is a Coxeter group, and

$$W_L = W_Q \rtimes (L/Q), \quad \text{where } L/Q \text{ is finite and abelian.}$$

The *length* $\ell(w)$ of $w \in W_Q$ is the minimum $\ell \geq 0$ such that w can be written as a product of ℓ generators in S . The length of $w \in W_L$ is defined by $\ell(w) = \ell(w')$, where $w = w'\gamma$ with $w' \in W_Q$ and $\gamma \in L/Q$. Thus elements of L/Q have length zero.

Write $R = R_1 \cup R_2 \cup R_3$ with

$$\begin{aligned} R_1 &= \{\alpha \in R \mid \alpha/2 \notin R \text{ and } 2\alpha \notin R\}, & R_2 &= \{\alpha \in R \mid 2\alpha \in R\}, \\ R_3 &= \{\alpha \in R \mid \alpha/2 \in R\}. \end{aligned}$$

These sets are pairwise disjoint, and if R is reduced then $R_1 = R$ and $R_2 = R_3 = \emptyset$. Define

$$R_0 = R_1 \cup R_2.$$

The *inversion set* of $w \in W$ is $R(w) = \{\alpha \in R_0^+ \mid w^{-1}\alpha \in -R_0^+\}$. By [3, VI, §1], we have

$$R(w) = \{\alpha_{i_1}, s_{i_1}\alpha_{i_2}, \dots, s_{i_1} \cdots s_{i_{\ell-1}}\alpha_{i_\ell}\} \quad \text{whenever } w = s_{i_1} \cdots s_{i_\ell} \text{ is reduced.} \quad (2.1)$$

For each rank $n \geq 1$ there is exactly one irreducible non-reduced root system (up to isomorphism). This is the BC_n system, and it can be realised in \mathbb{R}^n by

$$R = \pm\{e_i - e_j, e_i + e_j, e_k, 2e_k \mid 1 \leq i < j \leq n \text{ and } 1 \leq k \leq n\},$$

where the simple roots are $\alpha_i = e_i - e_{i+1}$ for $1 \leq i < n$ and $\alpha_n = e_n$. The coroot lattice is spanned by $\alpha_1^\vee, \dots, \alpha_{n-1}^\vee, \alpha_n^\vee/2$, and we have $P = Q$. Then R_0 is a root system of type B_n with simple roots $\alpha_1, \dots, \alpha_n$. We will always use the above conventions for indexing the simple roots of BC_n root systems, and more generally we will adopt standard Bourbaki conventions [3] for the irreducible root systems.

2.2 Parameter systems

Let $q_0, q_1, \dots, q_n \in \mathbb{C}^\times$ be such that $q_i = q_j$ whenever s_i and s_j are conjugate in W_Q . We call the sequence (q_i) a *parameter system*. By [3, IV, §5, No. 5, Prop. 5], the product

$$q_w = q_{i_1} \cdots q_{i_\ell} \quad (\text{where } w = s_{i_1} \cdots s_{i_\ell} \in W_Q \text{ is reduced})$$

does not depend on the particular reduced expression for w . If $\alpha \in W_0\alpha_i \cap W_0\alpha_j$ then s_i and s_j are conjugate in W_0 , and hence for $\alpha \in R_0$ we define

$$q_\alpha = q_i \quad \text{if } \alpha \in W_0\alpha_i.$$

Let $\mathbb{C}[L] = \mathbb{C}\text{-span}\{x^\lambda \mid \lambda \in L\}$ be the group algebra of L , with the group operation written multiplicatively as $x^\lambda x^\mu = x^{\lambda+\mu}$. In the field of fractions of $\mathbb{C}[L]$, let (for $\alpha \in R_0$)

$$c_\alpha(x) = \begin{cases} \frac{1 - q_\alpha^{-1}x^{-\alpha^\vee}}{1 - x^{-\alpha^\vee}} & \text{if } \alpha \in R_1, \\ \frac{(1 - q_0^{-1/2}q_n^{-1/2}x^{-\alpha^\vee/2})(1 + q_0^{1/2}q_n^{-1/2}x^{-\alpha^\vee/2})}{1 - x^{-\alpha^\vee}} & \text{if } \alpha \in R_2. \end{cases}$$

(Note that if $\alpha \in R_2$ then $2\alpha \in R$, and so $(2\alpha)^\vee = \alpha^\vee/2$ is in L .) Choose relatively prime elements $n_\alpha(x)$ and $d_\alpha(x)$ in $\mathbb{C}[L]$ so that

$$c_\alpha(x) = \frac{n_\alpha(x)}{d_\alpha(x)}.$$

For example, $\alpha \in R_2$ and $q_n^{1/2} = q_0^{1/2}$ then $n_\alpha(x) = 1 - q_n^{-1}x^{-\alpha^\vee/2}$ and $d_\alpha(x) = 1 - x^{-\alpha^\vee/2}$.

Similarly, let

$$c'_\alpha(x) = \begin{cases} \frac{1-q_\alpha^{-1}}{1-x^{-\alpha^\vee}} & \text{if } \alpha \in R_1, \\ \frac{1-q_n^{-1}+q_n^{-1/2}(q_0^{1/2}-q_0^{-1/2})x^{-\alpha^\vee/2}}{1-x^{-\alpha^\vee}} & \text{if } \alpha \in R_2. \end{cases}$$

Then, with $d_\alpha(x)$ as above,

$$c'_\alpha(x) = \frac{n'_\alpha(x)}{d_\alpha(x)}$$

for some $n'_\alpha(x) \in \mathbb{C}[L]$ with $n'_\alpha(x)$ and $d_\alpha(x)$ relatively prime.

Let

$$c(x) = \prod_{\alpha \in R_0^+} c_\alpha(x), \quad n(x) = \prod_{\alpha \in R_0^+} n_\alpha(x), \quad d(x) = \prod_{\alpha \in R_0^+} d_\alpha(x).$$

The expression $c(x) = n(x)/d(x)$ is the *Macdonald c-function*. We write $c_i(x), c'_i(x), n_i(x), n'_i(x)$, and $d_i(x)$ for $n_i(x)$ for $c_{\alpha_i}(x), c'_{\alpha_i}(x), n_{\alpha_i}(x), n'_{\alpha_i}(x)$, and $d_{\alpha_i}(x)$, respectively.

2.3 Affine Hecke algebras

Standard references for affine Hecke algebras include [14, 15] and [17]. With the above notation, the *affine Hecke algebra* \mathcal{H}_L with parameters q_0, \dots, q_n is the algebra over \mathbb{C} with vector space basis $\{T_w \mid w \in W_L\}$ and relations

$$T_u T_v = T_{uv} \quad \text{if } \ell(uv) = \ell(u) + \ell(v),$$

$$T_i^2 = 1 + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}})T_i \quad \text{for all } i = 0, 1, \dots, n,$$

where we write $T_i = T_{s_i}$.

The above presentation is the *Coxeter presentation* of \mathcal{H}_L . There is a second important presentation which exploits the semidirect product structure $W_L = L \rtimes W_0$. This is the *Bernstein presentation*, given by (2.2)–(2.5) below. Each $\lambda \in L$ can be written as $\lambda = \mu - \nu$ with $\mu, \nu \in L \cap P^+$, and we define

$$x^\lambda = T_{t_\mu} T_{t_\nu}^{-1}.$$

It is not hard to see that this is well defined, and in particular $x^\lambda = T_{t_\lambda}$ if λ is dominant.

It can be shown [15] that \mathcal{H}_L has vector space basis $\{T_w x^\lambda \mid \lambda \in L, w \in W_0\}$ and relations

$$T_i^2 = 1 + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}})T_i \quad \text{for } i = 1, \dots, n, \tag{2.2}$$

$$T_i T_j T_i \cdots = T_j T_i T_j \cdots \quad (m_{ij} \text{ factors}) \quad \text{for } 1 \leq i < j \leq n, \tag{2.3}$$

$$x^\lambda x^\mu = x^{\lambda+\mu} \quad \text{for all } \lambda, \mu \in L, \tag{2.4}$$

$$T_i x^\lambda - x^{s_i \lambda} T_i = q_i^{\frac{1}{2}} c'_i(x) (x^\lambda - x^{s_i \lambda}) \quad \text{for } 1 \leq i \leq n \text{ and } \lambda \in L. \tag{2.5}$$

Thus we see a copy of the group algebra $\mathbb{C}[L]$ of the lattice L inside of \mathcal{H}_L . The relation (2.5) is the *Bernstein relation*. Since $s_i \lambda = \lambda - \langle \lambda, \alpha_i \rangle \alpha_i^\vee$ and $\langle \lambda, \alpha_i \rangle \in \mathbb{Z}$ the “fraction” $c'_i(x) (x^\lambda - x^{s_i \lambda})$ that appears in the Bernstein relation is actually an element of $\mathbb{C}[L]$.

It is well known (see, for example, [15, (4.2.10)]) that the centre of \mathcal{H}_L is

$$\mathbb{C}[L]^{W_0} = \{f \in \mathbb{C}[L] \mid w \cdot f = f \text{ for all } w \in W_0\}.$$

This has powerful implications for the representation theory of \mathcal{H}_L . For example, it forces the irreducible representations to be finite dimensional (since the centre acts on irreducible representations by scalars, and it is evident that \mathcal{H}_L is finite dimensional over $\mathbb{C}[L]^{W_0}$).

It is natural to seek modifications τ_w of T_w which satisfy a “simplified Bernstein relation”

$$\tau_w x^\lambda = x^{w\lambda} \tau_w \quad \text{for all } w \in W_0 \text{ and } \lambda \in L. \tag{2.6}$$

Define elements $\tau_1, \dots, \tau_n \in \mathcal{H}_L$ by

$$\tau_i = d_i(x) T_i - q_i^{\frac{1}{2}} n'_i(x).$$

The Bernstein relation gives $\tau_i x^\lambda = x^{s_i \lambda} \tau_i$, and it can be shown (see [23, Proposition 2.7], for example) that for $w \in W_0$ the product

$$\tau_w = \tau_{i_1} \cdots \tau_{i_\ell} \quad \text{is independent of the choice of reduced expression } w = s_{i_1} \cdots s_{i_\ell}.$$

Thus the elements $\tau_w, w \in W_0$, satisfy (2.6), and a direct calculation gives the useful formula

$$\tau_i^2 = q_i n_i(x) n_i(x^{-1}) \in \mathbb{C}[L]. \tag{2.7}$$

2.4 Harmonic analysis for the affine Hecke algebra

Suppose now that $q_0, q_1, \dots, q_n > 1$. Define an involution $*$ on \mathcal{H}_L and the *canonical trace functional* $\text{Tr} : \mathcal{H}_L \rightarrow \mathbb{C}$ by

$$\left(\sum_{w \in W_L} c_w T_w \right)^* = \sum_{w \in W_L} \overline{c_w} T_{w^{-1}} \quad \text{and} \quad \text{Tr} \left(\sum_{w \in W_L} c_w T_w \right) = c_1.$$

An induction on $\ell(v)$ shows that $\text{Tr}(T_u T_v^*) = \delta_{u,v}$ for all $u, v \in W_L$, and so

$$\text{Tr}(h_1 h_2) = \text{Tr}(h_2 h_1) \quad \text{for all } h_1, h_2 \in \mathcal{H}_L.$$

Thus $\langle h_1, h_2 \rangle = \text{Tr}(h_1 h_2^*)$ defines an Hermitian inner product on \mathcal{H}_L . Let $\|h\|_2 = \sqrt{\langle h, h \rangle}$. The algebra \mathcal{H}_L acts on itself by left multiplication, and the corresponding operator norm is

$$\|h\| = \sup \{ \|hx\|_2 : x \in \mathcal{H}_L, \|x\|_2 \leq 1 \}.$$

Let $\overline{\mathcal{H}_L}$ denote the completion of \mathcal{H}_L with respect to this norm. Thus $\overline{\mathcal{H}_L}$ is a non-commutative C^* -algebra. This algebra is ‘liminal’. Even better, all irreducible representations of $\overline{\mathcal{H}_L}$ are finite dimensional, and so by [7, §8.8] there exists a probability measure μ such that

$$\text{Tr}(h) = \int_{\text{spec}(\overline{\mathcal{H}_L})} \chi_\pi(h) d\mu(\pi) \quad \text{for all } h \in \overline{\mathcal{H}_L}. \tag{2.8}$$

The measure μ is the *Plancherel measure*. Only those representations of \mathcal{H}_L which extend to the completion $\overline{\mathcal{H}_L}$ appear in the Plancherel Theorem. It is known [19, Corollary 6.2] that these are the *tempered* representations of \mathcal{H}_L (see [19, §2.7] for the definition).

If $t \in \text{Hom}(L, \mathbb{C}^\times)$ we write $t^\lambda = t(\lambda)$. The Weyl group W_0 acts on $\text{Hom}(L, \mathbb{C}^\times)$ by the formula $(wt)^\lambda = t^{w^{-1}\lambda}$. Following [18], define a function $G_t : \mathcal{H}_L \rightarrow \mathbb{C}$ by

$$G_t(h) = \sum_{\mu \in L} t^{-\mu} \text{Tr}(x^\mu h) \tag{2.9}$$

whenever the series converges. From [18] we have the following (see also [19, (3.9)]).

Theorem 2.1 *The series $G_t(h)$ is absolutely convergent for all $h \in \mathcal{H}_L$ whenever $|t^{\alpha_i^\vee}| < q_i^{-1}$ for $(R, i) \neq (BC_n, n)$ and $|t^{\alpha_n^\vee}| < q_0^{-1}q_n^{-1}$ for $(R, i) = (BC_n, n)$. Moreover,*

$$G_t(h) = \frac{g_t(h)}{q_{w_0}c(t)c(t^{-1})d(t)}, \tag{2.10}$$

where for each fixed h the function $g_t(h)$ has a analytic continuation in the t -variable to $\text{Hom}(L, \mathbb{C}^\times)$. Moreover, $g_t(h)$ satisfies

1. $g_t(h)$ is a polynomial in $\{t^\lambda \mid \lambda \in L\}$ (for fixed $h \in \mathcal{H}_L$),
2. $g_t(1) = d(t)$ for all $t \in \text{Hom}(L, \mathbb{C}^\times)$, and
3. $g_t(x^\lambda h x^\mu) = t^{\lambda+\mu} g_t(h)$ for all $\lambda, \mu \in L$ and all $h \in \mathcal{H}_L$.

Remark 2.2 (a) Note that $t^\lambda g_t(\tau_w) = g_t(\tau_w x^\lambda) = g_t(x^{w\lambda} \tau_w) = t^{w\lambda} g_t(\tau_w)$, and so if $wt \neq t$ then

$$g_t(\tau_w x^\lambda) = \delta_{w,1} t^\lambda d(t).$$

Then, by condition 1 in the theorem, this formula holds for all $t \in \text{Hom}(L, \mathbb{C}^\times)$.

(b) The three conditions in the theorem completely determine $g_t(h)$. For example, consider the \tilde{A}_2 case. It is sufficient to calculate $g_t(T_w)$ for each $w \in W_0$, because $g_t(T_w x^\lambda) = t^\lambda g_t(T_w)$. Write $Q = q^{\frac{1}{2}} - q^{-\frac{1}{2}}$. Applying g_t to the Bernstein relation $T_1 x^{\alpha_1^\vee} = x^{-\alpha_1^\vee} T_1 + Q(1 + x^{\alpha_1^\vee})$ gives

$$g_t(T_1) = Q(1 - t^{-\alpha_1^\vee})^{-1} d(t) = Q(1 - t^{-\alpha_2^\vee})(1 - t^{-\alpha_1^\vee - \alpha_2^\vee}).$$

Similarly, $g_t(T_2) = Q(1 - t^{-\alpha_1^\vee})(1 - t^{-\alpha_1^\vee - \alpha_2^\vee})$, $g_t(T_1 T_2) = g_t(T_2 T_1) = Q^2(1 - t^{-\alpha_1^\vee - \alpha_2^\vee})$, and $g_t(T_1 T_2 T_1) = Q^3 + Q(1 - t^{-\alpha_1^\vee})(1 - t^{-\alpha_2^\vee})$, making (2.10) completely explicit in type \tilde{A}_2 .

Let

$$f_i(h) = \frac{g_i(h)}{d(t)}. \tag{2.11}$$

Note that $f_i(h)$ may have poles at points where $t^{\alpha \vee} = 1$ for some $\alpha \in R_0$. Fix a \mathbb{Z} -basis $\lambda_1, \dots, \lambda_n$ of L . From (2.9) and (2.10) we have

$$\text{Tr}(h) = \frac{1}{q_{w_0}} \int_{a_1 \mathbb{T}} \cdots \int_{a_n \mathbb{T}} \frac{f_t(h)}{c(t)c(t^{-1})} dt_1 \cdots dt_n, \tag{2.12}$$

where $t_i = t^{\lambda_i}$, dt_i is Haar measure on the circle group \mathbb{T} , and where $a_1, \dots, a_n > 0$ are chosen such that if $t \in \text{Hom}(L, \mathbb{C}^\times)$ with $|t^{\lambda_i}| = a_i$ for each i then $|t^{\alpha_i \vee}| < q_i^{-1}$ (if $(R, i) \neq (BC_n, n)$) and $|t^{\alpha_n \vee}| < q_0^{-1} q_n^{-1}$ (if $(R, i) = (BC_n, n)$). Formula (2.12) appears in [19, Theorem 3.7], and is the starting point of the Plancherel Theorem.

3 Representations of affine Hecke algebras

Let M be a finite dimensional \mathcal{H}_L -module. For each $t \in \text{Hom}(L, \mathbb{C}^\times)$ let

$$M_t = \{v \in M \mid x^\lambda \cdot v = t^\lambda v \text{ for all } \lambda \in L\},$$

$$M_t^{\text{gen}} = \{v \in M \mid \text{for each } \lambda \in L \text{ there is a } k > 0 \text{ such that } (x^\lambda - t^\lambda)^k \cdot v = 0\}$$

be the t -weight space and the generalised t -weight space, respectively. Each finite dimensional \mathcal{H}_L -module M decomposes into a direct sum of generalised t -weight spaces

$$M = \bigoplus_{t \in \text{supp}(M)} M_t^{\text{gen}},$$

where $\text{supp}(M) = \{t \in \text{Hom}(L, \mathbb{C}^\times) \mid M_t^{\text{gen}} \neq 0\}$ is the support of M .

A finite dimensional \mathcal{H}_L -module M is calibrated if

$$M_t^{\text{gen}} = M_t \text{ for all } t \in \text{supp}(M).$$

(In the literature, this is also referred to as *tame*). The main purpose of the first half of this paper is to construct calibrated irreducible representations of general affine Hecke algebras. We suspect that a complete classification of calibrated representations along the lines of the 1-parameter case is possible (perhaps with some restrictions like $W_t = W_{(t)}$), although such a classification requires detailed information about the representation theory of rank 2 (multi-parameter) affine Hecke algebras, and this would take us beyond the scope of the present paper (the rank 2 representation theory for the 1-parameter case is treated in [25]). Thus the focus of this paper is on construction, and the question of classification will be pursued in later work.

The elements $\tau_i \in \mathcal{H}_L$, considered as operators on a representation, are often called *intertwining operators* because of the following fundamental and important fact.

Lemma 3.1 *Let $1 \leq i \leq n$. Let M be a finite dimensional \mathcal{H}_L -module, and suppose that $t \in \text{supp}(M)$. Then as operators,*

$$\tau_i : M_i^{\text{gen}} \rightarrow M_{s_i t}^{\text{gen}} \quad \text{and} \quad \tau_i : M_{s_i t}^{\text{gen}} \rightarrow M_t^{\text{gen}}.$$

Moreover, $n_i(t)n_i(t^{-1}) \neq 0$ if and only if each operator is bijective.

Proof Let $m \in M_i^{\text{gen}}$. By (2.6), we compute

$$(x^\lambda - (s_i t)^\lambda)^k \tau_i \cdot m = \tau_i (x^{s_i \lambda} - t^{s_i \lambda})^k \cdot m = 0$$

for sufficiently large k . Thus $\tau_i \cdot m \in M_{s_i t}^{\text{gen}}$. For the final claim, note that by (2.7) the operator $\tau_i^2 : M_i^{\text{gen}} \rightarrow M_t^{\text{gen}}$ is given by $\tau_i^2 \cdot m = q_i n_i(t) n_i(t^{-1}) m$. □

By Schur’s Lemma (see [29]), the centre $\mathbb{C}[L]^{W_0}$ of \mathcal{H}_L acts on an irreducible module M by scalars. It follows that there exists $t \in \text{Hom}(L, \mathbb{C}^\times)$ such that

$$p(x) \cdot v = p(t)v \quad \text{for all } p(x) \in \mathbb{C}[L]^{W_0} \text{ and all } v \in M.$$

The element t is only defined up to W_0 orbits. The orbit $W_0 t$ is called the *central character* of M , although as is customary we will usually refer to any $t' \in W_0 t$ as ‘the’ central character of M . A central character $t \in \text{Hom}(L, \mathbb{C}^\times)$ is called *regular* if $t^{\alpha^\vee} \neq 1$ for all $\alpha \in R_0$.

3.1 Principal series representations

The large commutative subalgebra $\mathbb{C}[L]$ of \mathcal{H}_L can be used to construct finite dimensional representations of \mathcal{H}_L . The *principal series representation* with *central character* $t \in \text{Hom}(L, \mathbb{C}^\times)$ is

$$M(t) = \text{Ind}_{\mathbb{C}[L]}^{\mathcal{H}_L}(\mathbb{C}v_t) = \mathcal{H}_L \bigotimes_{\mathbb{C}[L]} (\mathbb{C}v_t),$$

where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[L]$ with $x^\lambda \cdot v_t = t^\lambda v_t$ for all $\lambda \in L$. It is clear that this representation is $|W_0|$ -dimensional, and that $\{T_w \otimes v_t \mid w \in W_0\}$ is a basis of $M(t)$.

For $t \in \text{Hom}(L, \mathbb{C}^\times)$ define

$$N(t) = \{\alpha \in R_0^+ \mid n_\alpha(t)n_{-\alpha}(t) = 0\},$$

$$D(t) = \{\alpha \in R_0^+ \mid d_\alpha(t) = 0\}.$$

Note that $N(t)$ and $D(t)$ are closely related to the zeros of the numerator and denominator of the Macdonald c -function, respectively.

For $t \in \text{Hom}(L, \mathbb{C}^\times)$, let

$$W_t = \{w \in W_0 \mid wt = t\},$$

$$W_{(t)} = \{\{s_\alpha \mid \alpha \in D(t)\}\}.$$

Note that $W_{(t)}$ is a normal subgroup of W_t (since $ws_\alpha w^{-1} = s_{w\alpha}$). Moreover, if $L = P$ then necessarily $W_{(t)} = W_t$ (see [28, §4.2, 5.3]).

The following theorem of Kato [11, Theorem 2.2] is fundamental.

Theorem 3.2 *The module $M(t)$ is irreducible if and only if $N(t) = \emptyset$ and $W_t = W_{(t)}$.*

The fundamental importance of the principal series representations is highlighted by the following fact (see, for example, [23, Proposition 2.6]).

Proposition 3.3 *If M is an irreducible representation of \mathcal{H}_L with central character t then M is a quotient of $M(t)$. In particular, $\dim(M) \leq |W_0|$.*

3.2 A combinatorial construction of irreducible calibrated \mathcal{H}_L -modules

Following [23], the *calibration graph* of $t \in \text{Hom}(L, \mathbb{C}^\times)$ is the graph $\Gamma(t)$ with

$$\begin{aligned} &\text{vertex set } \{wt \mid w \in W_0\}, \quad \text{and} \\ &\text{edges } \{wt, s_i wt\} \quad \text{if and only if } \alpha_i \notin N(wt). \end{aligned}$$

For each $J \subseteq N(t)$ define

$$F_J(t) = \{w \in W_0 \mid R(w^{-1}) \cap N(t) = J \text{ and } R(w^{-1}) \cap D(t) = \emptyset\}.$$

By the argument in [23, Theorem 2.14], if $W_t = W_{(t)}$ then the connected components of $\Gamma(t)$ are precisely the sets

$$\{wt \mid w \in F_J(t)\} \quad \text{such that } J \subseteq N(t) \text{ and } F_J(t) \neq \emptyset. \tag{3.1}$$

Remark 3.4 We note that if $W_t = W_{(t)}$ then the geometric argument in [23, Theorem 2.14] also shows that if $w, v \in F_J(t)$, and if $wv^{-1} = s_{i_1} \cdots s_{i_\ell}$ is a reduced expression, then each element

$$w, s_{i_1} w, s_{i_2} s_{i_1} w, \dots, s_{i_\ell} \cdots s_{i_2} s_{i_1} w = v \quad \text{is in } F_J(t).$$

(Because the “smaller regions” in the proof of [23, Theorem 2.14] which correspond to the connected components of the calibration graph are convex in the sense that they are intersections of half spaces, and hence by [27, Proposition 2.8] all minimal length paths between w and v are contained in this region). Thus $F_J(t)$ is ‘geodesically closed’ in the (dual of the) underlying Coxeter complex.

Proposition 3.5 *If M is a finite dimensional \mathcal{H}_L -module then*

$$\dim(M_t^{\text{gen}}) = \dim(M_{t'}^{\text{gen}})$$

whenever t and t' are in the same connected component of the calibration graph $\Gamma(t)$.

Proof If $\alpha_i \notin N(t)$ then Lemma 3.1 gives $\dim(M_t^{\text{gen}}) = \dim(M_{s_i t}^{\text{gen}})$. Hence the result. \square

Let R_{ij} be the rank 2 subsystem of R generated by the simple roots α_i and α_j . That is, R_{ij} is the intersection of R with the \mathbb{Z} -span of $\{\alpha_i, \alpha_j\}$. We say that a weight $t \in \text{Hom}(L, \mathbb{C}^\times)$ is (i, j) -regular if $(wt)^{\alpha_i^\vee} \neq 1$ and $(wt)^{\alpha_j^\vee} \neq 1$ for all $w \in W_{ij} = \langle s_i, s_j \rangle$, and (i, j) -calibratable if one of the following conditions holds:

- (i) The weight t is (i, j) -regular.
- (ii) R_{ij} is of type C_2 (assume α_i short and α_j long) with
 - (a) $q_i = q_j$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee}) = (q_i, q_i^{-1})$ or (q_i^{-1}, q_i) , or
 - (b) $q_i = q_j^2$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee}) = (q_j^{-2}, q_j)$ or (q_j^2, q_j) .
- (iii) R_{ij} is of type G_2 (assume α_i short and α_j long) with
 - (a) $q_i = q_j$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee}) = (q_i^{-1}, q_i), (q_i, q_i^{-1}), (q_i^2, q_i^{-1}), (q_i^{-2}, q_i)$, or
 - (b) $q_i = q_j^3$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee}) = (q_j^3, q_j^{-1}), (q_j^{-3}, q_j), (q_j^{-3}, q_j^2), (q_j^3, q_j^{-2})$.
- (iv) R_{ij} is of type BC_2 (assume α_i middle-length and α_j short) with
 - (a) $q_i = q_0 q_j$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee/2}) = (q_0 q_j, q_0^{-1/2} q_j^{-1/2}), (q_0^{-1} q_j^{-1}, q_0^{1/2} q_j^{1/2})$, or
 - (b) $q_i = q_0 q_j^{-1}$ or $q_0^{-1} q_j$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee/2}) = (q_0^{-1} q_j, -q_0^{1/2} q_j^{-1/2}), (q_0 q_j^{-1}, -q_0^{-1/2} q_j^{1/2})$, or
 - (c) $q_i = q_0^{1/2} q_j^{1/2}$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee/2}) = (q_i^{-1}, q_i), (q_i, q_i^{-1})$, or
 - (d) $q_i = q_0^{-1/2} q_j^{1/2}$ or $q_0^{1/2} q_j^{-1/2}$ and $(t^{\alpha_i^\vee}, t^{\alpha_j^\vee/2}) = (q_i^{-1}, -q_i), (q_i, -q_i^{-1})$.

Conditions (i), (ii)(a) and (iii)(a) are equivalent to Ram’s definition of calibratable in the 1-parameter case. Note that if R_{ij} is of type BC_2 then the underlying root system R is necessarily non-reduced, and hence is of type BC_n , and so $1 \leq i \leq n - 1$ and $j = n$ (since α_j is assumed to be short).

In the following theorem, we construct a class of irreducible calibrated \mathcal{H}_L -modules.

Theorem 3.6 *Let $t \in \text{Hom}(L, \mathbb{C}^\times)$, and let $J \subseteq N(t)$. Suppose that $F_J(t) \neq \emptyset$ and that each wt with $w \in F_J(t)$ is (i, j) -calibratable for each pair (α_i, α_j) of simple roots of R . Let $M_J(t)$ be the vector space over \mathbb{C} with basis $\{e_{wt} \mid w \in F_J(t)\}$, and define linear operators \tilde{T}_i ($i = 1, \dots, n$), \tilde{x}^λ ($\lambda \in L$), on $M_J(t)$ by linearly extending the formulae*

$$\tilde{x}^\lambda e_{wt} = (wt)^\lambda e_{wt} \quad \lambda \in L, \tag{3.2}$$

$$\tilde{T}_i e_{wt} = q_i^{\frac{1}{2}} c'_i(wt) e_{wt} + q_i^{\frac{1}{2}} c_i(wt) e_{s_i wt} \quad 1 \leq i \leq n, \tag{3.3}$$

with the convention that $e_{vt} = 0$ if $v \notin F_J(t)$. Then the map $\mathcal{H}_L \rightarrow \text{End}(M_J(t))$ with $T_i \mapsto \tilde{T}_i$ and $x^\lambda \mapsto \tilde{x}^\lambda$ defines an irreducible calibrated representation of \mathcal{H}_L .

Proof (a) We check that the operators \tilde{T}_i and \tilde{x}^λ satisfy the Bernstein relation. We have

$$\begin{aligned}
 (\tilde{T}_i \tilde{x}^\lambda - \tilde{x}^{s_i \lambda} \tilde{T}_i) e_{wt} &= ((wt)^\lambda - \tilde{x}^{s_i \lambda}) \tilde{T}_i e_{wt} \\
 &= ((wt)^\lambda - x^{s_i \lambda}) (q_i^{\frac{1}{2}} c'_i(wt) e_{wt} + q_i^{\frac{1}{2}} c_i(wt) e_{s_i wt}) \\
 &= ((wt)^\lambda - (wt)^{s_i \lambda}) q_i^{\frac{1}{2}} c'_i(wt) e_{wt} = q_i^{\frac{1}{2}} c'_i(\tilde{x}) (\tilde{x}^\lambda - \tilde{x}^{s_i \lambda}) e_{wt}.
 \end{aligned}$$

(b) We now check that the operators \tilde{T}_i satisfy the quadratic relation $\tilde{T}_i^2 = 1 + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}}) \tilde{T}_i$.

$$\begin{aligned}
 \tilde{T}_i^2 e_{wt} &= \tilde{T}_i (q_i^{\frac{1}{2}} c'_i(wt) e_{wt} + q_i^{\frac{1}{2}} c_i(wt) e_{s_i wt}) \\
 &= q_i (c'_i(wt)^2 + c_i(wt) c_i(s_i wt)) e_{wt} + q_i c_i(wt) (c'_i(wt) + c'_i(s_i wt)) e_{s_i wt} \\
 &= (1 + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}}) q_i^{\frac{1}{2}} c'_i(wt)) e_{wt} + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}}) q_i^{\frac{1}{2}} c_i(wt) e_{s_i wt} \\
 &= (1 + (q_i^{\frac{1}{2}} - q_i^{-\frac{1}{2}}) \tilde{T}_i) e_{wt}.
 \end{aligned}$$

(c) We verify the braid relation $\dots \tilde{T}_i \tilde{T}_j \tilde{T}_i = \dots \tilde{T}_j \tilde{T}_i \tilde{T}_j$ (m_{ij} factors). Fix $w \in F_J(t)$. Suppose first that wt is (i, j) -regular. Let $v \in W_{ij}$. If $e_{vwt} \neq 0$ then (3.3) gives

$$(\tilde{T}_i - q_i^{\frac{1}{2}} c'_i(vwt)) e_{vwt} = q_i^{\frac{1}{2}} c_i(vwt) e_{s_i vwt}, \tag{3.4}$$

and, by Remark 3.4, this formula is also true when $e_{vwt} = 0$ and $\ell(s_i v) > \ell(v)$.

Consider the product (well defined by (i, j) -regularity)

$$\begin{aligned}
 A_{ij}(wt) &= \dots (\tilde{T}_i - q_i^{\frac{1}{2}} c'_i(s_j s_i wt)) (\tilde{T}_j - q_j^{\frac{1}{2}} c'_j(s_i wt)) (\tilde{T}_i - q_i^{\frac{1}{2}} c'_i(wt)) \\
 &\quad (m_{ij} \text{ factors}).
 \end{aligned}$$

Let v_0 be the longest element of W_{ij} . Repeatedly using (3.4) and $c_\alpha(vwt) = c_{v^{-1}\alpha}(wt)$ gives

$$A_{ij}(wt) e_{wt} = q_{v_0}^{\frac{1}{2}} [c_{\alpha_i}(wt) c_{s_i \alpha_j}(wt) c_{s_i s_j \alpha_i}(wt) c_{s_i s_j s_j \alpha_j}(wt) \dots] e_{v_0 wt}.$$

By (2.1), we have $\{\alpha_i, s_i \alpha_j, s_i s_j \alpha_i, \dots\} = \{\alpha_j, s_j \alpha_i, s_j s_i \alpha_j, \dots\}$ and so $A_{ji}(wt) e_{wt} = A_{ij}(wt) e_{wt}$. Each $v \in W_{ij} \setminus \{v_0\}$ has a unique expression as a product of simple generators, and so for $v < v_0$ we may unambiguously define operators $\tilde{T}_v = \tilde{T}_{i_1} \tilde{T}_{i_2} \dots \tilde{T}_{i_\ell}$ where $v = s_{i_1} s_{i_2} \dots s_{i_\ell}$ is the unique reduced expression for $v \in W_{ij}$. Expanding $A_{ij}(wt)$ and $A_{ji}(wt)$ and using the already verified quadratic relation for \tilde{T}_i and \tilde{T}_j , we see that there are rational functions $p_v(wt)$ and $q_v(wt)$ in wt such that

$$\begin{aligned}
 A_{ij}(wt) e_{wt} &= \dots \tilde{T}_i \tilde{T}_j \tilde{T}_i e_{wt} + \sum_{v < v_0} p_v(wt) \tilde{T}_v e_{wt}, \\
 A_{ji}(wt) e_{wt} &= \dots \tilde{T}_j \tilde{T}_i \tilde{T}_j e_{wt} + \sum_{v < v_0} q_v(wt) \tilde{T}_v e_{wt}.
 \end{aligned}$$

One now shows that $p_v(wt) = q_v(wt)$ for all $v < v_0$. This is achieved exactly as in [23, Proposition 2.7] by using the action of the τ -operators on principal series

representations, and we omit the details. Thus the braid relation, in the (i, j) -regular case, holds.

We now verify the braid relation in the case where wt is (i, j) -calibratable but not (i, j) -regular. Consider the $R_{ij} = C_2$ case with $q_i = q_j$ and $(wt)^{\alpha_i^\vee} = q_i$ and $(wt)^{\alpha_j^\vee} = q_i^{-1}$. By (3.1), we have $F_J(t) = \{wt\}$, and so the braid relation is trivially satisfied (as $M_J(t)$ is 1-dimensional). All other C_2 cases are similar. In the G_2 case with $q_i = q_j^3$ and $(wt)^{\alpha_i^\vee} = q_j^3$ and $(wt)^{\alpha_j^\vee} = q_j^{-2}$, by (3.1) we compute $F_J(t) = \{w, s_j w\}$, and a direct calculation gives

$$\begin{aligned} \tilde{T}_i e_{wt} &= q_i^{\frac{1}{2}} e_{wt}, & \tilde{T}_j e_{wt} &= \frac{1}{q_j + 1} (-q_j^{-\frac{1}{2}} e_{wt} + q_j^{\frac{1}{2}} e_{s_j wt}), \\ \tilde{T}_i e_{s_j wt} &= -q_i^{-\frac{1}{2}} e_{s_j wt}, & \tilde{T}_j e_{s_j wt} &= q_j^{\frac{1}{2}} \left(\frac{1 - q_j^{-3}}{1 - q_j^{-2}} e_{wt} + \frac{1}{1 + q_j^{-1}} e_{s_j wt} \right). \end{aligned}$$

The braid relation follows by direct calculation. The remaining G_2 cases are similar (or trivial). Finally, in all BC_2 cases we have $F_J(t) = \{wt\}$ and so the braid relation is trivially satisfied.

To conclude the proof, we show that the module $M_J(t)$ is irreducible and calibrated. By the construction, the generalised weight spaces of $M_J(t)$ are $M_J(t)_{wt}$, with $w \in F_J(t)$, and each generalised weight space has dimension 1. So $M_J(t)$ is calibrated. Furthermore, it follows that if M is a proper submodule of $M_J(t)$ then there is $w, w' \in F_J(t)$ with $wt \neq w't$ such that $M_{wt} \neq 0$ and $M_{w't} = 0$, contradicting Proposition 3.5. Thus $M_J(t)$ is irreducible. □

We note the following subtle point: In Theorem 3.6, the basis of $M_J(t)$ is indexed by the set $\{wt \mid w \in F_J(t)\}$, while in the construction [24, Theorem 3.5] the basis is indexed by $F_J(t)$. The reason for this refinement is that we work with general lattices $Q \subseteq L \subseteq P$, while in [23, 24] the lattice $L = P$ is specified. See Examples 1 and 2 below.

Remark 3.7 Recently [5], Ram’s construction has been applied to study the representation theory of 1-parameter rank 2 affine Hecke algebras with q a root of the Poincaré polynomial, and analogously the above construction could be applied to the study of such representations in the multi-parameter case.

3.3 Examples

Let us give some concrete examples of the construction from Theorem 3.6. Most of these examples will arise in the Plancherel Theorems in the later parts of this paper (see Sect. 4). Of interest, we see in the third and fourth examples that some non-calibrated modules (in the single parameter case) can be constructed from calibrated modules (of multi-parameter algebras) by making an appropriate change of basis and taking a limit.

Example 1 Let \mathcal{H} be a \tilde{C}_2 Hecke algebra with $L = P$ and with parameters q_1 and q_2 (see Sect. 4.5). Let $t \in \text{Hom}(P, \mathbb{C}^\times)$ be the character with $t^{\omega_1} = -q_1^{-1}$ and




Fig. 1 Calibration graphs for Examples 1 and 2

$t^{\omega_2} = q_1^{-1/2}$, so that $t^{\alpha_1^\vee} = q_1^{-1}$ and $t^{\alpha_2^\vee} = -1$. Thus $N(t)^\vee = \{\alpha_1^\vee, \alpha_1^\vee + 2\alpha_2^\vee\}$ and $D(t) = \emptyset$. Thus there are 4 choices for subsets $J \subseteq N(t)$. Let $J_1^\vee = \emptyset$, $J_2^\vee = \{\alpha_1^\vee\}$, $J_3^\vee = \{\alpha_1^\vee + 2\alpha_2^\vee\}$, and $J_4^\vee = \{\alpha_1^\vee, \alpha_1^\vee + 2\alpha_2^\vee\}$. We compute $F_{J_1}(t) = \{1, s_2\}$, $F_{J_2}(t) = \{s_1, s_2s_1\}$, $F_{J_3}(t) = \{s_1s_2, s_2s_1s_2\}$, and $F_{J_4}(t) = \{s_1s_2s_1, s_1s_2s_1s_2\}$, and so the calibration graph is as in Fig. 1(a). Thus by Theorem 3.6 and Proposition 3.3, there are 4 irreducible modules with central character t , each with dimension 2. For example, the matrices for the module $M_{J_1}(t)$ with respect to the basis $\{e_t, e_{s_2t}\}$ are $\pi(T_1) = -q_1^{-1/2}I$, $\pi(x^{\omega_1}) = -q_1^{-1}I$, and

$$\pi(T_2) = \frac{q_2^{1/2}}{2} \begin{pmatrix} 1 - q_2^{-1} & 1 + q_2^{-1} \\ 1 + q_2^{-1} & 1 - q_2^{-1} \end{pmatrix}, \quad \pi(x^{\omega_2}) = \text{diag}(q_1^{-1/2}, -q_1^{-1/2}).$$

Note that $\pi(x^{\alpha_1^\vee}) = q_1^{-1}I$ and $\pi(x^{\alpha_2^\vee}) = -I$, and it follows that the restriction $\pi|_{\mathcal{H}_Q}$ is not irreducible (indeed, $\pi|_{\mathcal{H}_Q}$ is the direct sum of the representations π^4 and π^5 from Sect. 4.4). This does not contradict the irreducibility statement of Theorem 3.6, because the calibration graph changes if we use the lattice Q instead of P (see Example 2).

Example 2 Now let \mathcal{H} be a \tilde{C}_2 Hecke algebra with $L = Q$. Let $t \in \text{Hom}(Q, \mathbb{C}^\times)$ be the character with $t^{\alpha_1^\vee} = q_1^{-1}$ and $t^{\alpha_2^\vee} = -1$ (note the similarity to Example 1). Then $N(t)$ and $D(t)$ are as in Example 1. Let J_1, J_2, J_3, J_4 be as in Example 1, and then the sets $F_{J_i}(t)$ are as computed in Example 1. However, now $s_2t = t$, and so the calibration graph is as shown in Fig. 1(b). Thus Theorem 3.6 constructs 2 irreducible 1-dimensional modules, and 1 irreducible 2-dimensional module with central character t .

Example 3 Let \mathcal{H} be a \tilde{G}_2 Hecke algebra with $L = Q = P$ and with parameters q_1 and q_2 (see Sect. 4.6). Let $t \in \text{Hom}(Q, \mathbb{C}^\times)$ be the character with $t^{\alpha_1^\vee} = q_1$ and $t^{\alpha_2^\vee} = q_1^{-1/2}q_2^{1/2}$. If $q_1 \neq q_2$ and $q_1 \neq q_2^3$ then this character is regular, and we compute $N(t)^\vee = \{\alpha_1^\vee, \alpha_1^\vee + 2\alpha_2^\vee\}$. Thus there are 4 choices for $J \subseteq N(t)$, and the connected components of the calibration graph are given by $\{wt \mid w \in F_J(t)\}$ for these choices of J . Consider the case $J^\vee = \{\alpha_1^\vee + 2\alpha_2^\vee\}$. We compute $F_J(t) =$

$\{s_2s_1s_2s_1, s_1s_2s_1s_2s_1, s_2s_1s_2s_1s_2s_1\}$. The matrices for $\pi = M_J(t)$ are

$$\pi(T_1) = q_1^{\frac{1}{2}} \begin{pmatrix} \frac{1-q_1^{-1}}{1-q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}}} & \frac{1-q_1^{-\frac{3}{2}}q_2^{\frac{3}{2}}}{1-q_1^{-\frac{1}{2}}q_2^{\frac{3}{2}}} & 0 \\ \frac{1-q_1^{-\frac{1}{2}}q_2^{-\frac{3}{2}}}{1-q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}}} & \frac{1-q_1^{-1}}{1-q_1^{-\frac{1}{2}}q_2^{\frac{3}{2}}} & 0 \\ 0 & 0 & -q_1^{-1} \end{pmatrix},$$

$$\pi(T_2) = q_2^{\frac{1}{2}} \begin{pmatrix} -q_2^{-1} & 0 & 0 \\ 0 & \frac{1-q_2^{-1}}{1-q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}}} & \frac{1-q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}}}{1-q_1^{-\frac{1}{2}}q_2^{\frac{1}{2}}} \\ 0 & \frac{1-q_1^{-\frac{1}{2}}q_2^{-\frac{1}{2}}}{1-q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}}} & \frac{1-q_2^{-1}}{1-q_1^{-\frac{1}{2}}q_2^{\frac{1}{2}}} \end{pmatrix},$$

$$\pi(x^{\alpha_1^\vee}) = \text{diag}(q_1^{-\frac{1}{2}}q_2^{\frac{3}{2}}, q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}}, q_1^{-1}), \quad \pi(x^{\alpha_2^\vee}) = \text{diag}(q_2^{-1}, q_1^{-\frac{1}{2}}q_2^{\frac{1}{2}}, q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}}).$$

The construction breaks down when $q_1 = q_2$ or when $q_1 = q_2^3$. These cases can be dealt with by a suitable change of basis in the module $M_J(t)$. Let

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}} \\ 0 & -1 & 1 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & -q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}} & 0 \\ -1 & 1 & 0 \\ 0 & 0 & 1 - q_1^{\frac{1}{2}}q_2^{-\frac{3}{2}} \end{pmatrix}.$$

After conjugating each representing matrix by A (resp., B) it is observed that the resulting matrices are defined at $q_1 = q_2$ (resp., $q_1 = q_2^3$). Setting $q_1 = q_2 = q$ (resp., $q_1 = q^3$ with $q_2 = q$) gives a (non-calibrated) irreducible representation of the algebra $\mathcal{H}(q, q)$ (resp., the algebra $\mathcal{H}(q, q^3)$). For example, the matrices in the $q_1 = q_2 = q$ case become

$$\pi(T_1) = q^{\frac{1}{2}} \begin{pmatrix} 1 & \frac{3}{q-1} & \frac{3}{q-1} \\ \frac{q+1}{q} & \frac{2q+1}{q(q-1)} & \frac{3}{q-1} \\ -\frac{q+1}{q} & -\frac{3}{q-1} & -\frac{4q-1}{q(q-1)} \end{pmatrix},$$

$$\pi(T_2) = q^{\frac{1}{2}} \begin{pmatrix} -q^{-1} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -2q^{-1} & -q^{-1} \end{pmatrix},$$

$$\pi(x^{\alpha_1^\vee}) = \begin{pmatrix} q & 0 & 0 \\ 0 & -2q^{-1} & -3q^{-1} \\ 0 & 3q^{-1} & 4q^{-1} \end{pmatrix}, \quad \pi(x^{\alpha_2^\vee}) = \begin{pmatrix} q^{-1} & 0 & 0 \\ 0 & 3 & 2 \\ 0 & -2 & -1 \end{pmatrix}.$$

Example 4 Let \mathcal{H} be a \tilde{C}_2 affine Hecke algebra with either $L = Q$ or $L = P$ and with parameters q_1 and q_2 (see Sects. 4.4 and 4.5). Let $t \in \text{Hom}(L, \mathbb{C}^\times)$ be a character with $t^{\alpha_1^\vee} = q_1^{-1}$ and $t^{\alpha_2^\vee} = q_2$. If $q_1 \neq q_2$ and $q_1 \neq q_2^2$ then the character t is regular, since $t^{\alpha_1^\vee + \alpha_2^\vee} = q_1^{-1}q_2$ and $t^{\alpha_1^\vee + 2\alpha_2^\vee} = q_1^{-1}q_2^2$. Thus we compute $N(t) = \{\alpha_1, \alpha_2\}$ and $D(t) = \emptyset$. There are 4 choices for $J \subseteq N(t)$, namely $J_1 = \emptyset$, $J_2 = \{\alpha_1\}$, $J_3 = \{\alpha_2\}$, and $J_4 = \{\alpha_1, \alpha_2\}$. We compute

$$\begin{aligned} F_{J_1}(t) &= \{1\}, & F_{J_2}(t) &= \{s_1, s_2s_1, s_1s_2s_1\}, \\ F_{J_3}(t) &= \{s_2, s_1s_2, s_2s_1s_2\}, & F_{J_4}(t) &= \{s_1s_2s_1s_2\}. \end{aligned}$$

The sets $\{wt \mid w \in F_{J_i}(t)\}$ with $i = 1, 2, 3, 4$ are the connected components of the calibration graph of t . Thus there are 4 irreducible modules $M_{J_i}(t)$ ($i = 1, 2, 3, 4$) with central character t , with dimensions 1, 3, 3, 1, respectively.

Consider the module $M_{J_3}(t)$ (this module will appear in the Plancherel Theorem for \tilde{C}_2). The matrices of $T_1, T_2, x^{\alpha_1^\vee}$ and $x^{\alpha_2^\vee}$ relative to the basis $e_{s_2t}, e_{s_1s_2t}, e_{s_2s_1s_2t}$ are

$$\begin{aligned} \pi(T_1) &= q_1^{\frac{1}{2}} \begin{pmatrix} \frac{1-q_1^{-1}}{1-q_1q_2^{-2}} & \frac{1-q_1^{-2}q_2^2}{1-q_1^{-1}q_2^2} & 0 \\ \frac{1-q_2^{-2}}{1-q_1q_2^{-2}} & \frac{1-q_1^{-1}}{1-q_1^{-1}q_2^2} & 0 \\ 0 & 0 & -q_1^{-1} \end{pmatrix}, \\ \pi(T_2) &= q_2^{\frac{1}{2}} \begin{pmatrix} -q_2^{-1} & 0 & 0 \\ 0 & \frac{1-q_2^{-1}}{1-q_1q_2^{-1}} & \frac{1-q_1^{-1}}{1-q_1^{-1}q_2} \\ 0 & \frac{1-q_1q_2^{-2}}{1-q_1q_2^{-1}} & \frac{1-q_2^{-1}}{1-q_1^{-1}q_2} \end{pmatrix}, \end{aligned}$$

$$\pi(x^{\alpha_1^\vee}) = \text{diag}(q_1^{-1}q_2^2, q_1q_2^{-2}, q_1^{-1}), \quad \pi(x^{\alpha_2^\vee}) = \text{diag}(q_2^{-1}, q_1^{-1}q_2, q_1q_2^{-1}).$$

If $L = P$ then $\omega_1 = \alpha_1^\vee + \alpha_2^\vee$ and $\omega_2 = \alpha_1^\vee/2 + \alpha_2^\vee$. Thus there are 2 characters $t \in \text{Hom}(P, \mathbb{C}^\times)$ with $t^{\alpha_1^\vee} = q_1^{-1}$ and $t^{\alpha_2^\vee} = q_2$, specifically $t^{\omega_1} = q_1^{-1}q_2$ and $t^{\omega_2} = \pm q_1^{-1/2}q_2$. The corresponding matrices for x^{ω_1} and x^{ω_2} are

$$\pi(x^{\omega_1}) = \text{diag}(q_1^{-1}q_2, q_2^{-1}, q_2^{-1}), \quad \pi(x^{\omega_2}) = \pm \text{diag}(q_1^{-1/2}, q_1^{-1/2}, q_1^{1/2}q_2^{-1}).$$

In the cases $q_1 = q_2$ or $q_1 = q_2^2$ an analogous computation to that in Example 2 can be used to construct (non-calibrated) irreducible representations of $\mathcal{H}(q, q)$ and $\mathcal{H}(q, q^2)$.

3.4 Characters

We conclude this section with some observations about characters that will be used for the Plancherel Theorems. Let $f_t(h)$ be as in (2.11).

Lemma 3.8 *Let π be an irreducible representation of \mathcal{H}_L with central character t , and suppose that the character χ of π satisfies*

$$\chi(\tau_w x^\lambda) = \delta_{w,1} \sum_{v \in W_0} k_v (vt)^\lambda \quad \text{for all } w \in W_0 \text{ and } \lambda \in L,$$

for some numbers $k_v \in \mathbb{C}$. Then, with f_t as in (2.11), if t is regular we have

$$\chi(h) = \sum_{v \in W_0} k_v f_{vt}(h) \quad \text{for all } h \in \mathcal{H}_L.$$

Proof Since t is regular, each $f_{vt}(h)$ with $v \in W_0$ and $h \in \mathcal{H}_L$ is defined. From Remark 2.2 and the hypothesis, we have $\chi(h) = \sum_{v \in W_0} k_v f_{vt}(h)$ for all $h \in \mathcal{H}'_L$, where \mathcal{H}'_L is the subalgebra of \mathcal{H}_L with basis $\{\tau_w x^\lambda \mid w \in W_0, \lambda \in L\}$. Let $\Delta(x) = \prod_{\alpha \in R_0} (1 - x^{-\alpha^\vee}) = d(x)d(x^{-1})$. An induction using the formula $(1 - x^{-\alpha^\vee})T_i = \tau_i + a_i(x)$ shows that $\Delta(x)^{\ell(w)}T_w \in \mathcal{H}'_L$ for all $w \in W_0$. Thus $\Delta(x)^{\ell(w)}T_w x^\lambda \in \mathcal{H}'_L$ for all $w \in W_0$ and $\lambda \in L$. Since $\Delta(x) \in \mathbb{C}[L]^{W_0}$ is central and χ is irreducible, we have

$$\begin{aligned} \Delta(t)^{\ell(w)}\chi(T_w x^\lambda) &= \chi(\Delta(x)^{\ell(w)}T_w x^\lambda) = \sum_{v \in W_0} k_v f_{vt}(\Delta(x)^{\ell(w)}T_w x^\lambda) \\ &= \Delta(t)^{\ell(w)} \sum_{v \in W_0} k_v f_{vt}(T_w x^\lambda). \end{aligned}$$

We can divide through by $\Delta(t)^{\ell(w)}$ since t is regular. □

Proposition 3.9 *Let χ_t be the character of the principal series representation $M(t)$ of \mathcal{H}_L with central character t . Then*

$$\chi_t(h) = \sum_{w \in W_0} f_{wt}(h) \quad \text{for all } h \in \mathcal{H}_L, \tag{3.5}$$

where the right hand side has an analytic continuation (for fixed $h \in \mathcal{H}_L$) to all $t \in \text{Hom}(L, \mathbb{C}^\times)$.

Proof Suppose first that $D(t) = \emptyset$ and that $M(t)$ is irreducible (see Theorem 3.2). Since $D(t) = \emptyset$ the module $M(t)$ has basis $\{\tau_w \otimes v_t \mid w \in W_0\}$. To see this note that if $w = s_{i_1} \cdots s_{i_\ell}$ is reduced then the Bernstein relation gives

$$\tau_w \otimes v_t = \left[\prod_{\alpha \in R(w^{-1})} (1 - t^{\alpha^\vee}) \right] (T_w \otimes v_t) + \text{lower terms},$$

where ‘lower terms’ is a linear combination of terms $T_v \otimes v_t$ with $v < w$ in Bruhat order. Thus if $D(t) = \emptyset$ then each basis element $T_w \otimes v_t$ of $M(t)$ can be written in terms of the elements $\{\tau_w \otimes v_t \mid w \in W_0\}$.

From (2.7) we see that the diagonal entries of the matrix for τ_w are all 0. The matrix for x^λ is diagonal with entries $(wt)^\lambda$ ($w \in W_0$) on the diagonal. Therefore,

$$\chi_t(\tau_w x^\lambda) = \delta_{w,1} \sum_{v \in W_0} (vt)^\lambda \quad \text{for all } w \in W_0 \text{ and } \lambda \in L.$$

Hence Lemma 3.8 gives (3.5).

The cases where $D(t) \neq \emptyset$ or $M(t)$ is not irreducible are obtained as follows. For fixed $h \in \mathcal{H}$ the character $\chi_t(h)$ is, by construction, a linear combination of $\{t^\lambda \mid \lambda \in L\}$ and is defined for all $t \in \text{Hom}(L, \mathbb{C}^\times)$. The right hand side of (3.5) is a rational function in t . Thus the singularities of this rational function are removable singularities (even though each individual summand may have singularities). \square

Proposition 3.10 *Suppose that t is a regular character. Let $J \subseteq N(t)$, and let $M_J(t)$ be the module constructed in Theorem 3.6. Then*

$$\chi(h) = \sum_{w \in F_J(t)} f_{wt}(h) \quad \text{for all } h \in \mathcal{H}.$$

Proof Since $\tau_i \cdot e_{wt} = q_i^{1/2} n_i(t) e_{s_i wt}$, we see that the diagonal entries of the matrix for τ_w are 0. Since the matrix representing x^λ is diagonal, it follows that

$$\chi(\tau_v x^\lambda) = \delta_{v,1} \sum_{w \in F_J(t)} (wt)^\lambda \quad \text{for all } v \in W_0 \text{ and } \lambda \in L,$$

and the result follows from Lemma 3.8. \square

Lemma 3.11 *Let π be a 1-dimensional representation of \mathcal{H}_L with regular central character t . Then*

$$\chi(h) = f_t(h) \quad \text{for all } h \in \mathcal{H}_L,$$

unless \mathcal{H}_L is of type \tilde{C}_n with $\pi(x^{\alpha_n^\vee}) = -1$. In this case there is a 1-dimensional representation π' defined by $\pi'(x^\lambda) = \pi(x^\lambda)$ for all $\lambda \in L$, $\pi'(T_i) = \pi(T_i)$ for all $i \neq n$, and $\pi'(T_n) = q_n^{1/2}$ (resp., $-q_n^{-1/2}$) if $\pi(T_n) = -q_n^{-1/2}$ (resp., $q_n^{1/2}$). Then

$$\frac{\chi(h) + \chi'(h)}{2} = f_t(h) \quad \text{for all } h \in \mathcal{H}_L.$$

Proof By direct analysis of the defining relations (2.2)–(2.5), one sees that the central character t of a 1-dimensional representation necessarily has $n_i(t)n_i(t^{-1}) = 0$, except in the \tilde{C}_n case with $\pi(x^{\alpha_n^\vee}) = -1$. Excluding this case for the moment, it follows from (2.7) that $\pi(\tau_i) = 0$ and hence $\pi(\tau_w x^\lambda) = \delta_{w,1} t^\lambda$ for all $w \in W_0$ and $\lambda \in L$. Since t is assumed to be regular, Lemma 3.8 gives $\chi(h) = f_t(h)$ for all $h \in \mathcal{H}_L$.

Now consider the \tilde{C}_2 case with $\pi(x^{\alpha_n^\vee}) = -1$. Let π' be the companion representation defined in the statement of the lemma. The proof of Lemma 3.8 applied to the representation $\pi \oplus \pi'$ proves the result. The fact that $\pi \oplus \pi'$ is not irreducible does not effect the proof of Lemma 3.8 because the centre $\mathbb{C}[L]^{W_0}$ of \mathcal{H}_L acts by the same scalar on each of π and π' . \square

Remark 3.12 In Proposition 3.10 and Lemma 3.11, we assumed that t is a regular central character. In general, these results are false for non-regular central characters, even if each term $f_t(h)$ is defined. For example, consider the \tilde{G}_2 case with $t \in \text{Hom}(Q, \mathbb{C}^\times)$ given by $t^{\alpha_1^\vee} = q_1, t^{\alpha_2^\vee} = q_2^{-1}$. If $q_1 \neq q_2$ and $q_1 \neq q_2^2$ and $q_1^2 \neq q_2^3$ and $q_1 \neq q_2^3$ then this central character is regular, and, by Lemma 3.11, we have $f_t(h) = \chi^4(h)$ for all $h \in \mathcal{H}$, where χ^4 is the 1-dimensional representation of \mathcal{H} listed in Sect. 4.6. Suppose that $q_1 = q_2 = q$. A calculation similar to Remark 2.2 shows that $f_t(h)$ is defined for all $h \in \mathcal{H}$, and that $f_t(T_1 T_2 T_1) = q^{1/2}$. But $\chi^4(T_1 T_2 T_1) = -q^{1/2}$.

4 The Plancherel Theorem

In this section, we state and prove the Plancherel Theorem for each irreducible affine Hecke algebra of rank 1 or rank 2. In each case, we give the generators and relations for the algebra, and construct the representations that appear in the Plancherel Theorem (see the Appendix for some explicit matrices). We then state the Plancherel Theorem, and give a proof starting from the trace generating function formula (2.12). The proof consists of performing a series of contour shifts and Proposition 3.9 to write (2.12) as

$$\text{Tr}(h) = \frac{1}{|W_0|q_{w_0}} \int_{\mathbb{T}^n} \frac{\chi_t(h)}{|c(t)|^2} dt + \text{lower terms}, \tag{4.1}$$

where the lower order terms are integrals over lower dimensional tori. Then the lower terms are matched up with lower dimensional representations of the Hecke algebra using Proposition 3.10 and Lemma 3.11.

Throughout this section, we assume that $q_0, q_1, \dots, q_n > 1$. The possible pairs (R, L) with R an irreducible rank 2 root system and L a \mathbb{Z} -lattice with $Q \subseteq L \subseteq P$ are $(R, L) = (A_2, Q), (A_2, P), (C_2, Q), (C_2, P), (G_2, Q),$ and (BC_2, Q) .

4.1 The rank 1 algebras

- (i) The $\tilde{A}_1(q), L = Q,$ algebra has generators $T = T_1$ and $x = x^{\alpha_1^\vee}$ with relations

$$T^2 = 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T, \quad Tx = x^{-1}T + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})(1 + x).$$

Let $\pi_t = \text{Ind}_{\text{Cl}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation with central character $t \in \mathbb{C}^\times$, where $x \cdot v_t = tv_t$. Let π be the 1-dimensional representation of \mathcal{H} with

$$\pi(T) = -q^{-\frac{1}{2}} \quad \text{and} \quad \pi(x) = q^{-1}.$$

Let χ_t be the character of π_t and let χ be the character of π .

- (ii) The $\tilde{A}_1(q), L = P,$ algebra has generators $T = T_1$ and $x = x^{\omega_1}$ with relations

$$T^2 = 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T, \quad Tx = x^{-1}T + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})x.$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[P]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation with central character $t \in \mathbb{C}^\times$, where $x \cdot v_t = tv_t$. Let π^1 and π^2 be the 1-dimensional representations of \mathcal{H} with

$$\begin{aligned} \pi^1(T) &= -q^{-\frac{1}{2}}, & \pi^1(x) &= q^{-\frac{1}{2}}, & \text{and } \pi^2(T) &= -q^{-\frac{1}{2}}, \\ \pi^2(x) &= -q^{-\frac{1}{2}}. \end{aligned}$$

Let χ_t, χ^1 , and χ^2 be the characters of π_t, π^1 , and π^2 , respectively.

- (iii) The $BC_1(q_0, q_1), L = Q$, algebra has generators $T = T_1, x = x^{\alpha_1^{1/2}}$ with relations

$$T^2 = 1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})T, \quad Tx = x^{-1}T + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})x + (q_0^{\frac{1}{2}} - q_0^{-\frac{1}{2}}).$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation with central character $t \in \mathbb{C}^\times$, where $x \cdot v_t = tv_t$. Let π^1, π^2 and π^3 be the 1-dimensional representations of \mathcal{H} with

$$\begin{aligned} \pi^1(T) &= -q_1^{-\frac{1}{2}}, & \pi^2(T) &= -q_1^{-\frac{1}{2}}, & \pi^3(T) &= q_1^{\frac{1}{2}}, \\ \pi^1(x) &= q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}, & \pi^2(x) &= -q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}, & \pi^3(x) &= -q_0^{-\frac{1}{2}}q_1^{\frac{1}{2}}. \end{aligned}$$

Let χ_t, χ^1, χ^2 , and χ^3 be the characters of π_t, π^1, π^2 , and π^3 , respectively.

Theorem 4.1 *Let $h \in \mathcal{H}$. In the cases (i), (ii) and (iii) above, we have, respectively:*

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{2q} \int_{\mathbb{T}} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{q-1}{q+1} \chi(h), \\ \text{Tr}(h) &= \frac{1}{2q} \int_{\mathbb{T}} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{q-1}{2(q+1)} (\chi^1(h) + \chi^2(h)), \\ \text{Tr}(h) &= \frac{1}{2q_1} \int_{\mathbb{T}} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{q_0q_1-1}{(q_0+1)(q_1+1)} \chi^1(h) + \frac{|q_0-q_1|}{(q_0+1)(q_1+1)} \\ &\quad \times \begin{cases} \chi^2(h) & \text{if } q_0 < q_1, \\ \chi^3(h) & \text{if } q_1 < q_0, \end{cases} \end{aligned}$$

where the c -functions are respectively

$$\begin{aligned} c(t) &= \frac{1-q^{-1}t^{-1}}{1-t^{-1}}, & c(t) &= \frac{1-q^{-1}t^{-2}}{1-t^{-2}}, \\ c(t) &= \frac{(1-q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}t^{-1})(1+q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}t^{-1})}{1-t^{-2}}. \end{aligned}$$

Proof Let us prove the $\tilde{BC}_1(q_0, q_1)$ case. If $q_0 = q_1$ there is some simplification, so suppose that $q_0 \neq q_1$. Write $g(t) = g_t(h)$ and $f(t) = f_t(h)$. From (2.12) we have

$$\text{Tr}(h) = \frac{1}{q_1} \int_{q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}a\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt,$$

where $0 < a < 1$. Note that the integrand has at most removable singularities on $t \in \mathbb{T}$, and that the poles of the integrand that lie between the contours $q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}a\mathbb{T}$ and \mathbb{T} are at $t = q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}, t = -q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}$ (in the case that $q_0 < q_1$) and $t = -q_0^{-\frac{1}{2}}q_1^{\frac{1}{2}}$ (in the case that $q_1 < q_0$). Computing residues (using $dt = \frac{1}{2\pi}d\theta = \frac{1}{2\pi i} \frac{dz}{z}$) gives

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{q_1} \int_{\mathbb{T}} \frac{f(t)}{|c(t)|^2} dt + \frac{q_0q_1 - 1}{(q_0 + 1)(q_1 + 1)} f(q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}) \\ &\quad + \frac{|q_0 - q_1|}{(q_0 + 1)(q_1 + 1)} \cdot \begin{cases} f(-q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}) & \text{if } q_0 < q_1, \\ f(-q_0^{-\frac{1}{2}}q_1^{\frac{1}{2}}) & \text{if } q_1 < q_0. \end{cases} \end{aligned}$$

Using Proposition 3.9, we have

$$\frac{1}{q_1} \int_{\mathbb{T}} \frac{f(t)}{|c(t)|^2} dt = \frac{1}{2q_1} \int_{\mathbb{T}} \frac{f(t) + f(t^{-1})}{|c(t)|^2} dt = \frac{1}{2q_1} \int_{\mathbb{T}} \frac{\chi_t(h)}{|c(t)|^2} dt,$$

and Lemma 3.11 gives $f(q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}) = \chi^1(h)$, $f(-q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}) = \chi^2(h)$, and $f(-q_0^{-\frac{1}{2}}q_1^{\frac{1}{2}}) = \chi^3(h)$. □

4.2 The $\tilde{A}_2(q)$ algebras with $L = Q$

This case is treated in [22], and so we will just state the result here. The coroot system is $R = \pm\{\alpha_1^\vee, \alpha_2^\vee, \alpha_1^\vee + \alpha_2^\vee\}$. The affine Hecke algebra has generators $T_1, T_2, x_1 = x^{\alpha_1^\vee}, x_2 = x^{\alpha_2^\vee}$ and relations

$$\begin{aligned} T_1^2 &= 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}T_1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})(1 + x_1), \\ T_1T_2T_1 &= T_2T_1T_2, & T_2^2 &= 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T_2, \\ T_2x_2 &= x_2^{-1}T_2 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})(1 + x_2), & x_1x_2 &= x_2x_1, \\ T_1x_2 &= x_1x_2T_1^{-1}, & T_2x_1 &= x_1x_2T_2^{-1}. \end{aligned}$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of the affine Hecke algebra \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[Q]$ with $x_1 \cdot v_t = t_1v_t$ and $x_2 \cdot v_t = t_2v_t$.

Let \mathcal{H}_1 be the subalgebra of \mathcal{H} generated by T_1, x_1 and x_2 . Let $s \in \mathbb{C}^\times$ and let $\mathbb{C}u_s$ be the 1-dimensional representation of \mathcal{H}_1 with

$$T_1 \cdot u_s = -q^{-\frac{1}{2}}u_s, \quad x_1 \cdot u_s = q^{-1}u_s, \quad x_2 \cdot u_s = q^{\frac{1}{2}}su_s.$$

Let $\pi_s^1 = \text{Ind}_{\mathcal{H}_1}^{\mathcal{H}}(\mathbb{C}u_s)$ be the induced representation of \mathcal{H} .

Let π^2 be the 1-dimensional representation of \mathcal{H} with

$$\pi^2(T_1) = -q^{-\frac{1}{2}}, \quad \pi^2(T_2) = -q^{-\frac{1}{2}}, \quad \pi^2(x_1) = q^{-1}, \quad \pi^2(x_2) = q^{-1}.$$

Let χ_t, χ_s^1 , and χ^2 be the characters of π_t, π_s^1 , and π^2 , respectively.

Theorem 4.2 *For all $h \in \mathcal{H}$ we have*

$$\text{Tr}(h) = \frac{1}{6q^3} \int_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{(q-1)^2}{q^2(q^2-1)} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds + \frac{(q-1)^3}{q^3-1} \chi^2(h),$$

where

$$c(t) = \frac{(1-q^{-1}t_1^{-1})(1-q^{-1}t_2^{-1})(1-q^{-1}t_1^{-1}t_2^{-1})}{(1-t_1^{-1})(1-t_2^{-1})(1-t_1^{-1}t_2^{-1})}, \quad c_1(s) = \frac{1-q^{-\frac{3}{2}}s^{-1}}{1-q^{\frac{1}{2}}s^{-1}}.$$

4.3 The $\tilde{A}_2(q)$ algebras with $L = P$

The root system is as in Sect. 4.2. The fundamental coweights are $\omega_1 = \frac{2}{3}\alpha_1^\vee + \frac{1}{3}\alpha_2^\vee$ and $\omega_2 = \frac{1}{3}\alpha_1^\vee + \frac{2}{3}\alpha_2^\vee$, and the coweight lattice is $P = \mathbb{Z}\omega_1 + \mathbb{Z}\omega_2$. The affine Hecke algebra is generated by $T_1, T_2, x_1 = x^{\omega_1}$ and $x_2 = x^{\omega_2}$ with relations

$$\begin{aligned} T_1^2 &= 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}x_2T_1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})x_1, \\ T_1T_2T_1 &= T_2T_1T_2, & T_2^2 &= 1 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})T_2, \\ T_2x_2 &= x_1x_2^{-1}T_2 + (q^{\frac{1}{2}} - q^{-\frac{1}{2}})x_2, & x_1x_2 &= x_2x_1, \\ T_1x_2 &= x_2T_1, & T_2x_1 &= x_1T_2. \end{aligned}$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[P]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of the affine Hecke algebra \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[P]$ with $x_1 \cdot v_t = t_1v_t$ and $x_2 \cdot v_t = t_2v_t$.

Let \mathcal{H}_1 be the subalgebra generated by T_1, x_1 and x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^1 = \mathcal{H} \otimes_{\mathcal{H}_1}(\mathbb{C}u_s)$ be the 3-dimensional representation of \mathcal{H} induced from the 1-dimensional representation $\mathbb{C}u_s$ of \mathcal{H}_1 given by

$$T_1 \cdot u_s = -q^{-\frac{1}{2}}u_s, \quad x_1 \cdot u_s = q^{-\frac{1}{2}}su_s, \quad x_2 \cdot u_s = s^2u_s.$$

The module π_s^1 has basis $\{1 \otimes u_s, T_2 \otimes u_s, T_1T_2 \otimes u_s\}$ and support $\text{supp } \pi_s^1 = \{t, s_2t, s_1s_2t\}$, where $t \in \text{Hom}(P, \mathbb{C}^\times)$ is the character with $(t^{\omega_1}, t^{\omega_2}) = (q^{-1/2}s, s^2)$. It is not hard to show (see the proof of Lemma 4.4) that the character of π_s^1 satisfies

$$\chi_s(h) = f_t(h) + f_{s_2t}(h) + f_{s_1s_2t}(h) \quad \text{for all } s \in \mathbb{C}^\times \text{ and all } h \in \mathcal{H}. \tag{4.2}$$

Let π^2, π^3 and π^4 be the 1-dimensional representations of \mathcal{H} given by (where $\omega = e^{2\pi i/3}$)

$$\begin{aligned} \pi^2(T_1) &= -q^{-\frac{1}{2}}, & \pi^2(T_2) &= -q^{-\frac{1}{2}}, & \pi^2(x_1) &= q^{-1}, & \pi^2(x_2) &= q^{-1}, \\ \pi^3(T_1) &= -q^{-\frac{1}{2}}, & \pi^3(T_2) &= -q^{-\frac{1}{2}}, & \pi^3(x_1) &= \omega q^{-1}, & \pi^3(x_2) &= \omega^{-1} q^{-1}, \\ \pi^4(T_1) &= -q^{-\frac{1}{2}}, & \pi^4(T_2) &= -q^{-\frac{1}{2}}, & \pi^4(x_1) &= \omega^{-1} q^{-1}, & \pi^4(x_2) &= \omega q^{-1}. \end{aligned}$$

Let $\chi_t, \chi_s^1, \chi^2, \chi^3$ and χ^4 be the characters of $\pi_t, \pi_s^1, \pi^2, \pi^3$, and π^4 , respectively.

Theorem 4.3 *For all $h \in \mathcal{H}$ we have*

$$\begin{aligned} \tau(h) &= \frac{1}{6q^3} \int_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{(q-1)^2}{q^2(q^2-1)} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds \\ &\quad + \frac{(q-1)^3}{3(q^3-1)} (\chi^2(h) + \chi^3(h) + \chi^4(h)), \end{aligned}$$

where

$$c(t) = \frac{(1 - q^{-1}t_1^{-2}t_2)(1 - q^{-1}t_1t_2^{-2})(1 - q^{-1}t_1^{-1}t_2^{-1})}{(1 - t_1^{-2}t_2)(1 - t_1t_2^{-2})(1 - t_1^{-1}t_2^{-1})}, \quad c_1(s) = \frac{1 - q^{-\frac{3}{2}}s^{-3}}{1 - q^{\frac{1}{2}}s^{-3}}.$$

Proof The series $G_t(h)$ converges whenever $|t^{\alpha_1^\vee}|, |t^{\alpha_2^\vee}| < q^{-1}$, and hence the series converges whenever $|t_1|, |t_2| < q^{-1}$, where $t_1 = t^{\omega_1}$ and $t_2 = t^{\omega_2}$. Fix $h \in \mathcal{H}$, and write $f_t(h) = f(t)$. Therefore,

$$\text{Tr}(h) = \frac{1}{q^3} \int_{q^{-1}a\mathbb{T}} \int_{q^{-1}b\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_1 dt_2,$$

where $0 < a, b < 1$. Fix a number $0 < c < 1$ very close to 1. Consider the inner integral. The t_1 -poles of the integrand lying between the contours $q^{-1}a\mathbb{T}$ and $c\mathbb{T}$ are at the points where $t_1^2 = q^{-1}t_2$. We compute the residues (using $dt_1 = \frac{1}{2\pi i} \frac{dz_1}{z_1}$) to be

$$\text{Res}_{t_1 = \pm q^{-1/2}t_2^{1/2}} \frac{f(t)}{c(t)c(t^{-1})} = -\frac{q(q-1)^2}{2(q^2-1)} \frac{f(\pm q^{-\frac{1}{2}}t_2^{1/2}, t_2)}{c_1(\mp t_2^{1/2})c_1(\mp t_2^{-1/2})}.$$

Using $\frac{1}{2} \int_{r\mathbb{T}} (f(t^{1/2}) + f(-t^{1/2})) dt = \int_{r^{1/2}\mathbb{T}} f(t) dt$ it follows that

$$\text{Tr}(h) = \frac{1}{q^3} \int_{q^{-1}a\mathbb{T}} \int_{c\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_1 dt_2 + \frac{(q-1)^2}{q^2(q^2-1)} \int_{q^{-\frac{1}{2}a} \frac{1}{2}\mathbb{T}} \frac{f(q^{-\frac{1}{2}}s, s^2)}{c_1(s)c_1(s^{-1})} ds.$$

Interchange the order of integration in the double integral. The t_2 -poles of the integrand between the contours $q^{-1}a\mathbb{T}$ to \mathbb{T} are at the points where $t_2^2 = q_1^{-1}t_1$ and where $t_2 = q^{-1}t_1^{-1}$. Computing residues gives

$$\text{Tr}(h) = \frac{1}{q^3} \int_{c\mathbb{T}} \int_{\mathbb{T}} \frac{f(t)}{|c(t)|^2} dt_2 dt_1 + \frac{(q-1)^2}{q^2(q^2-1)} (I_1 + I_2 + I_3),$$

where

$$I_1 = \int_{q^{-\frac{1}{2}a}c\mathbb{T}} \frac{f(q^{-\frac{1}{2}}s, s^2)}{c_1(s)c_1(s^{-1})} ds, \quad I_2 = \int_{c^{\frac{1}{2}}\mathbb{T}} \frac{f(s^2, q^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds,$$

$$I_3 = \int_{q^{\frac{1}{2}}c\mathbb{T}} \frac{f(q^{-\frac{1}{2}}s, q^{-\frac{1}{2}}s^{-1})}{c_1(s)c_1(s^{-1})} ds$$

and we have set $s = t_1^{1/2}$ in I_2 and $s = q^{\frac{1}{2}}t_1$ in I_3 . The t_1 -contour in the double integral can be shifted to \mathbb{T} without encountering any poles.

The plan is to shift each of the contours in I_1, I_2 and I_3 to the unit contour \mathbb{T} . However, we need to be careful with the possible singularities of $f(t)$. Therefore, we write $f(t) = g(t)/d(t)$, with $g(t)$ analytic. Then the integrands of the integrals I_1, I_2 and I_3 are

$$\frac{f(q^{-\frac{1}{2}}s, s^2)}{c_1(s)c_1(s^{-1})} = \frac{(1 - q^{\frac{1}{2}}s^3)g(q^{-\frac{1}{2}}s, s^2)}{(1 - s^{-2})(1 - q^{\frac{1}{2}}s^{-1})(1 - q^{-\frac{3}{2}}s^{-3})(1 - q^{-\frac{3}{2}}s^3)},$$

$$\frac{f(s^2, q^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} = \frac{(1 - q^{\frac{1}{2}}s^3)g(s^2, q^{-\frac{1}{2}}s)}{(1 - s^{-2})(1 - q^{\frac{1}{2}}s^{-1})(1 - q^{-\frac{3}{2}}s^{-3})(1 - q^{-\frac{3}{2}}s^3)},$$

$$\frac{f(q^{-\frac{1}{2}}s, q^{-\frac{1}{2}}s^{-1})}{c_1(s)c_1(s^{-1})} = \frac{(1 - q^{\frac{1}{2}}s^{-3})(1 - q^{\frac{1}{2}}s^3)g(q^{-\frac{1}{2}}s, q^{-\frac{1}{2}}s^{-1})}{(1 - q)(1 - q^{-\frac{3}{2}}s^{-3})(1 - q^{-\frac{3}{2}}s^3)(1 - q^{\frac{1}{2}}s^{-1})(1 - q^{\frac{1}{2}}s)}.$$

In particular, the integrands of I_1 and I_2 have singularities on \mathbb{T} . So instead we shift all contours to $c\mathbb{T}$. For the integrals I_2 and I_3 , we encounter no poles, and so the shift is for free. For the integral I_1 , we pick up simple residues at the points $s^3 = q^{-\frac{3}{2}}$, and computing residues gives

$$I_1 = \int_{c\mathbb{T}} \frac{f(q^{-\frac{1}{2}}s, s^2)}{c_1(s)c_1(s^{-1})} ds + \frac{q^2(q - 1)(q^2 - 1)}{3(q^3 - 1)} \times (f(q^{-1}, q^{-1}) + f(\omega q^{-1}, \omega^{-1}q^{-1}) + f(\omega^{-1}q^{-1}, \omega q^{-1})).$$

Therefore,

$$\text{Tr}(h) = \frac{1}{q^3} \int_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt$$

$$+ \frac{(q - 1)^2}{q^2(q^2 - 1)} \int_{c\mathbb{T}} \frac{f(q^{-\frac{1}{2}}s, s^2) + f(s^2, q^{-\frac{1}{2}}s) + f(q^{-\frac{1}{2}}s, q^{-\frac{1}{2}}s^{-1})}{c_1(s)c_1(s^{-1})} ds$$

$$+ \frac{(q - 1)^3}{3(q^3 - 1)} (f(q^{-1}, q^{-1}) + f(\omega q^{-1}, \omega^{-1}q^{-1}) + f(\omega^{-1}q^{-1}, \omega q^{-1})).$$

By (4.2), the numerator of the single integral is $\chi_s(h)$, and is therefore defined on \mathbb{T} and so the contour of the single integral can be shifted to \mathbb{T} . Proposition 3.9 deals with the double integral, and Lemma 3.11 deals with the 3 constant terms. □

4.4 The $\tilde{C}_2(q_1, q_2)$ algebras with $L = Q$

The dual root system is $R^\vee = \pm\{\alpha_1^\vee, \alpha_2^\vee, \alpha_1^\vee + \alpha_2^\vee, \alpha_1^\vee + 2\alpha_2^\vee\}$. Writing $x_1 = x^{\alpha_1^\vee}$ and $x_2 = x^{\alpha_2^\vee}$, the Hecke algebra has generators T_1, T_2, x_1, x_2 with relations

$$\begin{aligned} T_1^2 &= 1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}T_1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})(1 + x_1), \\ T_1T_2T_1T_2 &= T_2T_1T_2T_1, & T_2^2 &= 1 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})T_2, \\ T_2x_2 &= x_2^{-1}T_2 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})(1 + x_2), & x_1x_2 &= x_2x_1, \\ T_1x_2 &= x_1x_2T_1^{-1}, & T_2x_1 &= x_1x_2^2T_2^{-1} - (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})x_1x_2. \end{aligned}$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[Q]$ with $x_1 \cdot v_t = t_1 v_t$ and $x_2 \cdot v_t = t_2 v_t$.

Let \mathcal{H}_1 be the subalgebra generated by T_1, x_1, x_2 and let \mathcal{H}_2 be the subalgebra generated by T_2, x_1, x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^1 = \text{Ind}_{\mathcal{H}_1}^{\mathcal{H}}(\mathbb{C}u_s^1)$ and $\pi_s^2 = \text{Ind}_{\mathcal{H}_2}^{\mathcal{H}}(\mathbb{C}u_s^2)$ be the 4-dimensional representations induced from the 1-dimensional representation $\mathbb{C}u_s^1$ of \mathcal{H}_1 and the 1-dimensional representation $\mathbb{C}u_s^2$ of \mathcal{H}_2 given by

$$\begin{aligned} T_1 \cdot u_s^1 &= -q_1^{-\frac{1}{2}}u_s^1, & x_1 \cdot u_s^1 &= q_1^{-1}u_s^1, & x_2 \cdot u_s^1 &= q_1^{\frac{1}{2}}su_s^1, \\ T_2 \cdot u_s^2 &= -q_2^{-\frac{1}{2}}u_s^2, & x_1 \cdot u_s^2 &= q_2su_s^2, & x_2 \cdot u_s^2 &= q_2^{-1}u_s^2. \end{aligned}$$

Let π^j ($j = 3, 4, 5, 6, 7$) be the 1-dimensional representations of \mathcal{H} with

$$\begin{aligned} \pi^3(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^3(T_2) &= -q_2^{-\frac{1}{2}}, & \pi^3(x_1) &= q_1^{-1}, & \pi^3(x_2) &= q_2^{-1}, \\ \pi^4(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^4(T_2) &= -q_2^{-\frac{1}{2}}, & \pi^4(x_1) &= q_1^{-1}, & \pi^4(x_2) &= -1, \\ \pi^5(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^5(T_2) &= q_2^{\frac{1}{2}}, & \pi^5(x_1) &= q_1^{-1}, & \pi^5(x_2) &= -1, \\ \pi^6(T_1) &= q_1^{\frac{1}{2}}, & \pi^6(T_2) &= -q_2^{-\frac{1}{2}}, & \pi^6(x_1) &= q_1, & \pi^6(x_2) &= q_2^{-1}, \\ \pi^7(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^7(T_2) &= q_2^{\frac{1}{2}}, & \pi^7(x_1) &= q_1^{-1}, & \pi^7(x_2) &= q_2. \end{aligned}$$

Suppose that $q_1 \neq q_2$ and $q_1 \neq q_2^2$. Let $\pi^8 = M_J(t)$ be the representation with

$$(t^{\alpha_1^\vee}, t^{\alpha_2^\vee}) = (q_1^{-1}, q_2), \quad J^\vee = \{\alpha_2^\vee\}, \quad F_J(t) = \{s_2, s_1s_2, s_2s_1s_2\}$$

(since $q_1 \neq q_2$ and $q_1 \neq q_2^2$, we compute $N(t)^\vee = \{\alpha_1^\vee, \alpha_2^\vee\}$ and $D(t)^\vee = \emptyset$). The matrices for π^8 are given in Example 3 of Sect. 3.3.

Let $\chi_t, \chi_s^1, \chi_s^2$, and χ^j be the characters of π_t, π_s^1, π_s^2 , and π^j , respectively ($j = 3, \dots, 8$).

Lemma 4.4 *Let $t, u \in \text{Hom}(Q, \mathbb{C}^\times)$ be $(t^{\alpha_1^\vee}, t^{\alpha_2^\vee}) = (q_1^{-1}, q_1^{\frac{1}{2}}s)$ and $(u^{\alpha_1^\vee}, u^{\alpha_2^\vee}) = (q_2s, q_2^{-1})$ where $s \in \mathbb{C}^\times$. For all $h \in \mathcal{H}$ and all $s \in \mathbb{C}^\times$, we have*

$$\chi_s^1(h) = f_t(h) + f_{s_2t}(h) + f_{s_1s_2t}(h) + f_{s_2s_1s_2t}(h), \tag{4.3}$$

$$\chi_s^2(h) = f_u(h) + f_{s_1u}(h) + f_{s_2s_1u}(h) + f_{s_1s_2s_1u}(h). \tag{4.4}$$

Proof Let us prove (4.3) ((4.4) is similar). Suppose that $s \in \mathbb{C}^\times$ is not one of the isolated points of \mathbb{C}^\times which give $t^{\alpha^\vee} = 1$ for some $\alpha \in R_0^+$. Then π_s^1 is irreducible (for example, it can be constructed using Theorem 3.6 in these cases) and each $f_{vt}(h)$ is defined (for $v \in W_0$ and $h \in \mathcal{H}$). Moreover, π_s^1 has basis $\{1 \otimes u_s^1, \tau_2 \otimes u_s^1, \tau_1 \tau_2 \otimes u_s^1, \tau_2 \tau_1 \tau_2 \otimes u_s^1\}$ (this is proved in a similar way to the corresponding statement in the proof of Proposition 3.9).

The diagonal entries of each matrix $\pi_s^1(\tau_w)$ relative to this basis are 0. This is easily seen once it is observed that $\tau_1 \otimes u_s^1 = 0$ (which can be seen by direct calculation, or by (2.7)). Since

$$\pi_s^1(x^\lambda) = \text{diag}(t^\lambda, (s_2t)^\lambda, (s_1s_2t)^\lambda, (s_2s_1s_2t)^\lambda) \quad \text{for all } \lambda \in Q,$$

it follows that

$$\chi_s^1(\tau_w x^\lambda) = \delta_{w,1}(t^\lambda + (s_2t)^\lambda + (s_1s_2t)^\lambda + (s_2s_1s_2t)^\lambda) \quad \text{for all } w \in W_0 \text{ and } \lambda \in Q.$$

Thus Lemma 3.8 gives (4.3), provided s is not one of the isolated points of \mathbb{C}^\times that gives $t^{\alpha^\vee} = 1$ for some $\alpha \in R_0^+$. But by construction, $\chi_s^1(h)$ is a polynomial in s and s^{-1} (for fixed $h \in \mathcal{H}$) and the right-hand side of (4.3) is a rational function in s . Hence the result. □

Theorem 4.5 *For all $h \in \mathcal{H}$ we have*

$$\begin{aligned} \text{Tr}(h) = & \frac{1}{8q_1^2q_2^2} \iint_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{q_1 - 1}{2q_1q_2^2(q_1 + 1)} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds \\ & + \frac{q_2 - 1}{2q_1^2q_2(q_2 + 1)} \int_{\mathbb{T}} \frac{\chi_s^2(h)}{|c_2(s)|^2} ds \\ & + A\chi^3(h) + B(\chi^4(h) + \chi^5(h)) + |C| \times \begin{cases} \chi^6(h) & \text{if } q_1 < q_2, \\ \chi^8(h) & \text{if } q_2 < q_1 < q_2^2, \\ \chi^7(h) & \text{if } q_2^2 < q_1, \end{cases} \end{aligned}$$

where $c(t), c_1(s), c_2(s), A, B,$ and C are as in Appendix A.1. If $q_1 = q_2$ or $q_1 = q_2^2$ then the final term in the Plancherel Theorem is 0.

Proof The trace functional is given by

$$\text{Tr}(h) = \frac{1}{q_1^2q_2^2} \int_{q_1^{-1}a\mathbb{T}} \int_{q_2^{-1}b\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_2 dt_1, \tag{4.5}$$

where $0 < a, b < 1$ and where $f(t) = f_t(h)$. Choose a with $a < q_1q_2^{-1}$.

Step 1: Shifting the t_2 -contour. Let $0 < c < 1$ with $c^2 > q_1^{-1}$, $c > q_2^{-1}$, $c > q_1q_2^{-1}$ (if $q_1 < q_2$) and $c > q_1^{-1}q_2$ (if $q_2 < q_1$). We will shift the t_2 -contour from $q_2^{-1}b\mathbb{T}$ to $c\mathbb{T}$. The integrand has exactly one t_2 -pole between these contours, at $t_2 = q_2^{-1}$. Thus

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{q_1^2q_2^2} \int_{q_1^{-1}a\mathbb{T}} \int_{c\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_2 dt_1 + I_1, \quad \text{where} \\ I_1 &= -\frac{1}{q_1^2q_2^2} \int_{q_1^{-1}a\mathbb{T}} \text{Res}_{t_2=q_2^{-1}} \frac{f(t)}{c(t)c(t^{-1})} dt. \end{aligned}$$

Step 2: Shifting the t_1 -contour. Interchange the order of integration in the double integral. We will shift the t_1 -contour from $q_1^{-1}a\mathbb{T}$ to \mathbb{T} . By the conditions on a and c , the t_1 -poles of the integrand between these contours are at $t_1 = q_1^{-1}$, $t_1 = q_1^{-1}t_2^{-2}$, and $t_1 = q_2^{-1}t_2^{-1}$. Therefore,

$$\text{Tr}(h) = \frac{1}{q_1^2q_2^2} \int_{c\mathbb{T}} \int_{\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_1 dt_2 + I_1 + I_2 + I_3 + I_4,$$

where

$$I_j = -\frac{1}{q_1^2q_2^2} \int_{c\mathbb{T}} \text{Res}_{t_1=z_j} \frac{f(t)}{c(t)c(t^{-1})} dt_2 \quad \text{for } j = 2, 3, 4,$$

with $z_2 = q_1^{-1}$, $z_3 = q_1^{-1}t_2^{-2}$, and $z_4 = q_2^{-1}t_2^{-1}$. In the double integral, we may now revert back to the original order of integration, and shift the t_2 -contour to \mathbb{T} without encountering any poles.

Step 3: Shifting the contours in the integrals I_j . Straightforward calculations give

$$\begin{aligned} I_1 &= \frac{(q_2 - 1)^2}{q_1^2q_2(q_2^2 - 1)} \int_{q_1^{-1}q_2^{-1}a\mathbb{T}} \frac{f(q_2s, q_2^{-1})}{c_2(s)c_2(s^{-1})} ds, \\ I_2 &= \frac{(q_1 - 1)^2}{q_1q_2^2(q_1^2 - 1)} \int_{q_1^{-\frac{1}{2}}c\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds, \\ I_3 &= \frac{(q_1 - 1)^2}{q_1q_2^2(q_1^2 - 1)} \int_{q_1^{\frac{1}{2}}c\mathbb{T}} \frac{f(s^{-2}, q_1^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds, \\ I_4 &= \frac{(q_2 - 1)^2}{q_1^2q_2(q_2^2 - 1)} \int_{c\mathbb{T}} \frac{f(q_2^{-1}s^{-1}, s)}{c_2(s)c_2(s^{-1})} ds, \end{aligned}$$

where we have set $s = q_2^{-1}t_1$ in I_1 , $s = q_1^{-\frac{1}{2}}t_2$ in I_2 , $s = q_1^{\frac{1}{2}}t_2$ in I_3 , and $s = t_2$ in I_4 .

We now shift each contour to \mathbb{T} . As in the \tilde{A}_2 case, we need to be a little careful with possible singularities of $f(t)$. Thus we write $f(t) = g(t)/d(t)$. Then the

integrands of I_1, I_2, I_3 and I_4 are

$$\begin{aligned} \frac{f(q_2s, q_2^{-1})}{c_2(s)c_2(s^{-1})} &= \frac{q_2s(1-s)g(q_2s, q_2^{-1})}{(q_2-1)n_2(s)n_2(s^{-1})}, \\ \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} &= \frac{q_1^{\frac{1}{2}}s(1-s^2)g(q_1^{-1}, q_1^{\frac{1}{2}}s)}{(q_1-1)n_1(s)n_1(s^{-1})}, \\ \frac{f(s^{-2}, q_1^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} &= \frac{(1-s^{-2})g(s^{-2}, q_1^{-\frac{1}{2}}s)}{(1-q_1)n_1(s)n_1(s^{-1})}, \\ \frac{f(q_2^{-1}s^{-1}, s)}{c_2(s)c_2(s^{-1})} &= \frac{(1-s)g(q_2^{-1}s^{-1}, s)}{(1-q_2)n_2(s)n_2(s^{-1})}, \end{aligned}$$

where $n_1(s)$ and $n_2(s)$ are the numerators of $c_1(s)$ and $c_2(s)$. Each integrand is non-singular on \mathbb{T} (with removable singularities in the cases $q_1 = q_2$ or $q_1 = q_2^2$).

The poles of the integrand of I_1 which lie between the contours $q_1^{-1}q_2^{-1}a\mathbb{T}$ and \mathbb{T} are at $s = q_1^{-1}q_2^{-1}, s = q_1^{-1}q_2$ (if $q_2 < q_1$) and at $s = q_1q_2^{-1}$ (if $q_1 < q_2$). Calculating residues gives

$$\begin{aligned} I_1 &= \frac{(q_2-1)^2}{q_1^2q_2(q_2^2-1)} \int_{\mathbb{T}} \frac{f(q_2s, q_2^{-1})}{|c_2(s)|^2} ds + Af(q_1^{-1}, q_2^{-1}) \\ &+ C \times \begin{cases} f(q_1^{-1}q_2^2, q_2^{-1}) & \text{if } q_2 < q_1, \\ -f(q_1, q_2^{-1}) & \text{if } q_1 < q_2. \end{cases} \end{aligned}$$

The poles of the integrand of I_2 which lie between the contours $q_1^{-\frac{1}{2}}b\mathbb{T}$ and \mathbb{T} are at $s = -q_1^{-\frac{1}{2}}, s = q_1^{\frac{1}{2}}q_2^{-1}$ (if $q_2 < q_1 < q_2^2$), and $s = q_1^{-\frac{1}{2}}q_2$ (if $q_2^2 < q_1$). Calculating residues gives

$$\begin{aligned} I_2 &= \frac{(q_1-1)^2}{q_1q_2^2(q_1^2-1)} \int_{\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds + 2Bf(q_1^{-1}, -1) \\ &+ C \times \begin{cases} f(q_1^{-1}, q_1q_2^{-1}) & \text{if } q_2 < q_1 < q_2^2, \\ -f(q_1^{-1}, q_2) & \text{if } q_2^2 < q_1. \end{cases} \end{aligned}$$

The poles of the integrand of I_3 which lie between the contours $q_1^{\frac{1}{2}}c\mathbb{T}$ and \mathbb{T} are at $s = q_1^{-\frac{1}{2}}q_2$ (if $q_2 < q_1 < q_2^2$) and $s = q_1^{\frac{1}{2}}q_2^{-1}$ (if $q_2^2 < q_1$). Noting that \mathbb{T} is inside $q_1^{\frac{1}{2}}c\mathbb{T}$ gives

$$I_3 = \frac{(q_1-1)^2}{q_1q_2^2(q_1^2-1)} \int_{\mathbb{T}} \frac{f(s^{-2}, q_1^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds + C \times \begin{cases} f(q_1q_2^{-2}, q_1^{-1}q_2) & \text{if } q_2 < q_1 < q_2^2, \\ -f(q_1^{-1}q_2^2, q_2^{-1}) & \text{if } q_2^2 < q_1. \end{cases}$$

The integrand of I_4 has no poles between $c\mathbb{T}$ and \mathbb{T} . Therefore,

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{q_1^2 q_2^2} \iint_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt + \frac{(q_1 - 1)^2}{q_1 q_2^2 (q_1^2 - 1)} \int_{\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s) + f(s^{-2}, q_1^{-\frac{1}{2}}s)}{|c_1(s)|^2} ds \\ &+ \frac{(q_2 - 1)^2}{q_1^2 q_2 (q_2^2 - 1)} \int_{\mathbb{T}} \frac{f(q_2s, q_2^{-1}) + f(q_2^{-1}s^{-1}, s)}{|c_2(s)|^2} ds + Af(q_1^{-1}, q_2^{-1}) \\ &+ 2Bf(q_1^{-1}, -1) + |C| \\ &\times \begin{cases} f(q_1, q_2^{-1}) & \text{if } q_1 < q_2, \\ f(q_1^{-1}q_2^2, q_2^{-1}) + f(q_1^{-1}, q_1q_2^{-1}) + f(q_1q_2^{-2}, q_1^{-1}q_2) & \text{if } q_2 < q_1 < q_2^2, \\ f(q_1^{-1}, q_2) & \text{if } q_2^2 < q_1. \end{cases} \end{aligned}$$

Step 4: Matching with the representations. By Proposition 3.9, the double integral in the above formula is

$$\int_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt = \frac{1}{8} \int_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt.$$

The first single integral is

$$\begin{aligned} &\frac{1}{2} \int_{\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s) + f(s^{-2}, q_1^{-\frac{1}{2}}s) + f(q_1^{-1}, q_1^{\frac{1}{2}}s^{-1}) + f(s^2, q_1^{-\frac{1}{2}}s^{-1})}{|c_1(s)|^2} ds \\ &= \frac{1}{2} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds, \end{aligned}$$

where we have used Lemma 4.4. A similar analysis applies to the second single integral. Using Lemma 3.11, we have $f(q_1^{-1}, q_2^{-1}) = \chi^3(h)$ and $2f(q_1^{-1}, -1) = \chi^4(h) + \chi^5(h)$. Furthermore, for parameters $q_1 < q_2$ the central character $(t_1, t_2) = (q_1, q_2^{-1})$ is regular (for $t^{\alpha_1^\vee + \alpha_2^\vee} = q_1q_2^{-1} < 1$ and $t^{\alpha_1^\vee + 2\alpha_2^\vee} = q_1q_2^{-2} < 1$), and so Lemma 3.11 gives $f(q_1, q_2^{-1}) = \chi^6(h)$. Similarly, we have $f(q_1^{-1}, q_2) = \chi^7(h)$ for parameters $q_2^2 < q_1$. Finally, by Proposition 3.10, we have

$$f(q_1^{-1}q_2^2, q_2^{-1}) + f(q_1^{-1}, q_1q_2^{-1}) + f(q_1q_2^{-2}, q_1^{-1}q_2) = \chi^8(h)$$

for all parameters in the range $q_2 < q_1 < q_2^2$ (as the central character is regular). \square

4.5 The $\tilde{C}_2(q_1, q_2)$ algebras with $L = P$

The root system is as in Sect. 4.4, and the fundamental coweights are given by $\omega_1 = \alpha_1^\vee + \alpha_2^\vee$ and $\omega_2 = \frac{1}{2}\alpha_1^\vee + \alpha_2^\vee$. Writing $x_1 = x^{\omega_1}$ and $x_2 = x^{\omega_2}$, the Hecke algebra has presentation given by generators T_1, T_2, x_1, x_2 with relations

$$\begin{aligned} T_1^2 &= 1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}x_2^2T_1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})x_1, \\ T_1T_2T_1T_2 &= T_2T_1T_2T_1, & T_2^2 &= 1 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})T_2, \end{aligned}$$

$$T_2x_2 = x_1x_2^{-1}T_2 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})x_2, \quad x_1x_2 = x_2x_1,$$

$$T_1x_2 = x_2T_1, \quad T_2x_1 = x_1T_2.$$

The representation theory of \mathcal{H} is closely related to the representation theory of the Hecke algebra from Sect. 4.4.

Let $\pi_t = \text{Ind}_{\mathbb{C}[P]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[P]$ with $x_1 \cdot v_t = t_1v_t$ and $x_2 \cdot v_t = t_2v_t$.

Let \mathcal{H}_1 be the subalgebra generated by T_1, x_1, x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^\pm = \text{Ind}_{\mathcal{H}_1}^{\mathcal{H}}(\mathbb{C}u_s^\pm)$ be the 4-dimensional representations of \mathcal{H} induced from the representations $\mathbb{C}u_s^\pm$ of \mathcal{H}_1 with

$$T_1 \cdot u_s^\pm = -q_1^{-\frac{1}{2}}u_s^\pm, \quad x_1 \cdot u_s^\pm = q_1^{-\frac{1}{2}}su_s^\pm, \quad x_2 \cdot u_s^\pm = \pm su_s^\pm.$$

Let \mathcal{H}_2 be the subalgebra generated by T_2, x_1, x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^2 = \text{Ind}_{\mathcal{H}_2}^{\mathcal{H}}(\mathbb{C}u_s^2)$ be the 4-dimensional representation of \mathcal{H} induced from the representation $\mathbb{C}u_s^2$ of \mathcal{H}_2 with

$$T_2 \cdot u_s^2 = -q_2^{-\frac{1}{2}}u_s^2, \quad x_1 \cdot u_s^2 = s^2u_s^2, \quad x_2 \cdot u_s^2 = q_2^{-\frac{1}{2}}su_s^2.$$

Let π_\pm^j ($j = 3, 4, 5$) be the 1-dimensional representations

$$\begin{aligned} \pi_\pm^3(T_1) &= -q_1^{-\frac{1}{2}}, & \pi_\pm^3(T_2) &= -q_2^{-\frac{1}{2}}, & \pi_\pm^3(x_1) &= q_1^{-1}q_2^{-1}, \\ \pi_\pm^3(x_2) &= \pm q_1^{-\frac{1}{2}}q_2^{-1}, & \pi_\pm^4(T_1) &= q_1^{\frac{1}{2}}, & \pi_\pm^4(T_2) &= -q_2^{-\frac{1}{2}}, \\ \pi_\pm^4(x_1) &= q_1q_2^{-1}, & \pi_\pm^4(x_2) &= \pm q_1^{\frac{1}{2}}q_2^{-1}, & \pi_\pm^5(T_1) &= -q_1^{-\frac{1}{2}}, \\ \pi_\pm^5(T_2) &= q_2^{\frac{1}{2}}, & \pi_\pm^5(x_1) &= q_1^{-1}q_2, & \pi_\pm^5(x_2) &= \pm q_1^{-\frac{1}{2}}q_2. \end{aligned}$$

Let $\pi^6 = M_J(t)$ be the 2-dimensional representation with

$$(t^{\omega_1}, t^{\omega_2}) = (-q_1^{-1}, q_1^{-\frac{1}{2}}), \quad J^\vee = \emptyset, \quad F_J(t) = \{1, s_2\}$$

(we have $N(t)^\vee = \{\alpha_1^\vee, \alpha_2^\vee + 2\alpha_2^\vee\}$ and $D(t)^\vee = \emptyset$). Coincidentally, $\pi^6 \cong \text{Ind}_{\mathcal{H}_Q}^{\mathcal{H}}(\mathbb{C}u)$ where \mathcal{H}_Q is the algebra from Sect. 4.4 and where $\mathbb{C}u$ is the 1-dimensional representation of \mathcal{H}_Q with $T_1 \cdot u = -q_1^{-1/2}u$, $T_2 \cdot u = -q_2^{-1/2}u$, $x_1^\vee \cdot u = q_1^{-1}u$, and $x_2^\vee \cdot u = -u$. The matrices for π^6 are given in Example 1 of Sect. 3.3.

Suppose that $q_1 \neq q_2$ and $q_1 \neq q_2^2$. Let $\pi_\pm^7 = M_J(t_\pm)$ be the 3-dimensional representations with

$$(t_\pm^{\omega_1}, t_\pm^{\omega_2}) = (q_1^{-1}q_2, \pm q_1^{-\frac{1}{2}}q_2), \quad J^\vee = \{\alpha_2^\vee\}, \quad F_J(t_\pm) = \{s_2, s_1s_2, s_2s_1s_2\}$$

(we calculate $N(t_\pm)^\vee = \{\alpha_1^\vee, \alpha_2^\vee\}$ and $D(t)^\vee = \emptyset$).

Theorem 4.6 For all $h \in \mathcal{H}$ we have

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{8q_1^2q_2^2} \iint_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{(q_1 - 1)^2}{4q_1q_2^2(q_1^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^+(h) + \chi_s^-(h)}{|c_1(s)|^2} ds \\ &+ \frac{(q_2 - 1)^2}{2q_1^2q_2(q_2^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^2(h)}{|c_2(s)|^2} ds \\ &+ \frac{A}{2} (\chi_+^3(h) + \chi_-^3(h)) + B\chi^6(h) \\ &+ \frac{|C|}{2} \times \begin{cases} \chi_+^4(h) + \chi_-^4(h) & \text{if } q_1 < q_2, \\ \chi_+^7(h) + \chi_-^7(h) & \text{if } q_2 < q_1 < q_2^2, \\ \chi_+^5(h) + \chi_-^5(h) & \text{if } q_2^2 < q_1, \end{cases} \end{aligned}$$

where $c(t), c_1(s), c_2(s)$ are as in Appendix A.2 and A, B, C are as in Appendix A.1. If $q_1 = q_2$ or $q_1 = q_2^2$ then the final term in the Plancherel Theorem is 0.

Proof The series $G_t(h)$ converges for $|t^{\alpha_1^\vee}| < q_1^{-1}$ and $|t^{\alpha_2^\vee}| < q_2^{-1}$. Since $\alpha_1^\vee = 2\omega_1 - 2\omega_2$ and $\alpha_2^\vee = -\omega_1 + 2\omega_2$, the series converges whenever $|t_1^2t_2^{-2}| < q_1^{-1}$ and $|t_1^{-1}t_2^2| < q_2^{-1}$. Thus, writing $|t_1| = q_1^{-1}q_2^{-1}a$ and $|t_2| = q_1^{-1/2}q_2^{-1}b$, the series converges for $b^2 < a < b < 1$, and so

$$\text{Tr}(h) = \frac{1}{q_1^2q_2^2} \int_{q_1^{-1}q_2^{-1}a\mathbb{T}} \int_{q_1^{-1/2}q_2^{-1}b\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_2 dt_1.$$

From here one can either perform the contour shifts as in the $L = Q$ case, or change variables $t_1 = u_1u_2$ and $t_2^2 = u_1u_2^2$ to transform the above integral into $\frac{1}{2}$ times the $L = Q$ integral (4.5) with the numerator of its integrand replaced by $f^j(u_1, u_2) = f(u_1u_2, u_1^{1/2}u_2) + f(u_1u_2, -u_1^{1/2}u_2)$. We omit the details. \square

4.6 The $\tilde{G}_2(q_1, q_2)$ algebras with $L = Q$

The coroot system is $R^\vee = \pm\{\alpha_1^\vee, 2\alpha_1^\vee + 3\alpha_2^\vee, \alpha_1^\vee + 3\alpha_2^\vee, \alpha_2^\vee, \alpha_1^\vee + 2\alpha_2^\vee, \alpha_1^\vee + \alpha_2^\vee\}$, and the reflections s_1 and s_2 are given by $s_1(\alpha_2^\vee) = \alpha_1^\vee + \alpha_2^\vee$ and $s_2(\alpha_1^\vee) = \alpha_1^\vee + 3\alpha_2^\vee$. Writing $x_1 = x^{\alpha_1^\vee}$ and $x_2 = x^{\alpha_2^\vee}$, the Hecke algebra \mathcal{H} has generators T_1, T_2, x_1 and x_2 with relations

$$\begin{aligned} T_1^2 &= 1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}T_1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})(1 + x_1), \\ T_2^2 &= 1 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})T_2, & T_2x_2 &= x_2^{-1}T_2 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})(1 + x_2), \\ T_1T_2T_1T_2T_1T_2 &= T_2T_1T_2T_1T_2T_1, & T_2x_1 &= x_1x_2^3T_2^{-1} - (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})x_1x_2(1 + x_2), \\ x_1x_2 &= x_2x_1, & T_1x_2 &= x_1x_2T_1^{-1}. \end{aligned}$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[Q]$ with $x_1 \cdot v_t = t_1 v_t$ and $x_2 \cdot v_t = t_2 v_t$.

Let \mathcal{H}_1 be the subalgebra of \mathcal{H} generated by T_1, x_1, x_2 , and let \mathcal{H}_2 be the subalgebra generated by T_2, x_1, x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^1 = \text{Ind}_{\mathcal{H}_1}^{\mathcal{H}}(\mathbb{C}u_s^1)$ and $\pi_s^2 = \text{Ind}_{\mathcal{H}_2}^{\mathcal{H}}(\mathbb{C}u_s^2)$ be the 6-dimensional representations induced from the 1-dimensional representation $\mathbb{C}u_s^1$ of \mathcal{H}_1 and the 1-dimensional representation $\mathbb{C}u_s^2$ of \mathcal{H}_2 given by

$$\begin{aligned} T_1 \cdot u_s^1 &= -q_1^{-\frac{1}{2}} u_s^1, & x_1 \cdot u_s^1 &= q_1^{-1} u_s^1, & x_2 \cdot u_s^1 &= q_1^{\frac{1}{2}} s u_s^1, \\ T_2 \cdot u_s^2 &= -q_2^{-\frac{1}{2}} u_s^2, & x_1 \cdot u_s^2 &= q_2^{\frac{3}{2}} s u_s^2, & x_2 \cdot u_s^2 &= q_2^{-1} u_s^2. \end{aligned}$$

Let π^3, π^4 and π^5 be the 1-dimensional representations of \mathcal{H} with

$$\begin{aligned} \pi^3(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^3(T_2) &= -q_2^{-\frac{1}{2}}, & \pi^3(x_1) &= q_1^{-1}, & \pi^3(x_2) &= q_2^{-1}, \\ \pi^4(T_1) &= q_1^{\frac{1}{2}}, & \pi^4(T_2) &= -q_2^{-\frac{1}{2}}, & \pi^4(x_1) &= q_1, & \pi^4(x_2) &= q_2^{-1}, \\ \pi^5(T_1) &= -q_1^{-\frac{1}{2}}, & \pi^5(T_2) &= q_2^{\frac{1}{2}}, & \pi^5(x_1) &= q_1^{-1}, & \pi^5(x_2) &= q_2. \end{aligned}$$

Suppose that $q_1 \neq q_2, q_1 \neq q_2^2, q_1^2 \neq q_2^3, q_1 \neq q_2^3$. Let $\pi^6 = M_J(t)$ the 5-dimensional representation with

$$\begin{aligned} (t^{\alpha_1^\vee}, t^{\alpha_2^\vee}) &= (q_1^{-1}, q_2), & J^\vee &= \{\alpha_2^\vee\}, \\ F_J(t) &= \{s_2, s_1 s_2, s_2 s_1 s_2, s_1 s_2 s_1 s_2, s_2 s_1 s_2 s_1 s_2\}. \end{aligned}$$

Let $\pi_\pm^7 = M_J(t_\pm)$ be the 3-dimensional representations with

$$\begin{aligned} (t_\pm^{\alpha_1^\vee}, t_\pm^{\alpha_2^\vee}) &= (q_1, \pm q_1^{-\frac{1}{2}} q_2^{\frac{1}{2}}), & J^\vee &= \{\alpha_1^\vee + 2\alpha_2^\vee\}, \\ F_J(t_\pm) &= \{s_2 s_1 s_2 s_1, s_1 s_2 s_1 s_2 s_1, s_2 s_1 s_2 s_1 s_2 s_1\}, \end{aligned}$$

where we assume that $q_1 \neq q_2$ and $q_1 \neq q_2^3$ for $M_J(t_+)$. When $q_1 = q_2$ or $q_1 = q_2^3$, we define π_\pm^7 differently, as explained in Example 2 of Sect. 3.3.

Let $\pi^8 = M_J(t)$ be the 2-dimensional representation with

$$\begin{aligned} (t^{\alpha_1^\vee}, t^{\alpha_2^\vee}) &= (q_1, \omega), & J^\vee &= \{\alpha_1^\vee + 3\alpha_2^\vee\}, \\ F_J(t) &= \{s_1 s_2 s_1 s_2 s_1, s_2 s_1 s_2 s_1 s_2 s_1\}. \end{aligned}$$

Let $\chi_t, \chi_s^1, \chi_s^2, \chi^3, \chi^4, \chi^5, \chi^6, \chi_\pm^7$, and χ^8 be the characters of the above representations.

Theorem 4.7 *For all $h \in \mathcal{H}$ we have*

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{12q_1^3 q_2^3} \iint_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + \frac{(q_1 - 1)^2}{2q_1 q_2^3 (q_1^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds \\ &+ \frac{(q_2 - 1)^2}{2q_1^3 q_2^2 (q_2^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^2(h)}{|c_2(s)|^2} ds \end{aligned}$$

$$\begin{aligned}
 &+ A\chi^3(h) + B_+\chi_+^7(h) + B_-\chi_-^7(h) + C\chi^8(h) \\
 &+ |D| \times \begin{cases} \chi^4(h) & \text{if } q_1 < q_2^{3/2}, \\ \chi^6(h) & \text{if } q_2^{3/2} < q_1 < q_2^2, \\ \chi^5(h) & \text{if } q_2^2 < q_1, \end{cases}
 \end{aligned}$$

where $c(t), c_1(s), c_2(s), A, B_{\pm}, C, D$ are as in Appendix A.3. If $q_1 = q_2^{3/2}$ or $q_1 = q_2^2$ then the final term in the Plancherel Theorem is 0.

Proof Writing $f(t) = f_t(h)$, the trace functional is given by

$$\text{Tr}(h) = \frac{1}{q_1^3 q_2^3} \int_{q_1^{-1} a\mathbb{T}} \int_{q_2^{-1} b\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_2 dt_1$$

with $0 < a, b < 1$. Choose $0 < a, b < 1$ both very close to 0. Let $0 < c < 1$ be very close to 1. Consider the inner integral. The integrand has exactly one t_2 -pole between the contours $q_2^{-1}b\mathbb{T}$ and $c\mathbb{T}$, at $t_2 = q_2^{-1}$. Thus we can shift the t_2 -contour to $c\mathbb{T}$ at the cost of including this residue contribution. Now interchange the order of integration in the double integral. Since $|t_2| = c$, we see that the t_1 -poles of the integrand between the contours $q_1^{-1}a\mathbb{T}$ and \mathbb{T} are at the points where $t_1 = q_1^{-1}, t_1 = q_1^{-1}t_2^{-3}, t_1 = q_2^{-1}t_2^{-2}, t_1 = q_2^{-1}t_2^{-1}$ and $t_1^2 = q_1^{-1}t_2^{-3}$. After shifting the t_1 -contour to \mathbb{T} , we interchange the order of integration again, and since there are no t_2 -poles between $c\mathbb{T}$ and \mathbb{T} we shift the t_2 -contour to \mathbb{T} . Thus

$$\text{Tr}(h) = \frac{1}{q_1^3 q_2^3} \iint_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt + I_1 + I_2 + I_3 + I_4 + I_5 + I_6^+ + I_6^-,$$

where

$$\begin{aligned}
 I_1 &= -\frac{1}{q_1^3 q_2^3} \int_{q_1^{-1} a\mathbb{T}} \text{Res}_{t_2=z_1} \frac{f(t)}{c(t)c(t^{-1})} dt_1, & I_6^{\pm} &= -\frac{1}{q_1^3 q_2^3} \int_{c\mathbb{T}} \text{Res}_{t_1=\pm z_6} \frac{f(t)}{c(t)c(t^{-1})} dt_2, \\
 I_j &= -\frac{1}{q_1^3 q_2^3} \int_{c\mathbb{T}} \text{Res}_{t_1=z_j} \frac{f(t)}{c(t)c(t^{-1})} dt_2 \quad (j = 2, 3, 4, 5),
 \end{aligned}$$

where $z_1 = q_2^{-1}, z_2 = q_1^{-1}, z_3 = q_1^{-1}t_2^{-3}, z_4 = q_2^{-1}t_2^{-2}, z_5 = q_2^{-1}t_2^{-1}$, and $z_6 = q_1^{-\frac{1}{2}}t_2^{-\frac{3}{2}}$.

Use $\frac{1}{2} \int_{r\mathbb{T}} (f(t^{1/2}) + f(-t^{1/2})) dt = \int_{r^{1/2}\mathbb{T}} f(t) dt$ to write $I_6^+ + I_6^- = I_6$ as a single integral. Straightforward calculations give

$$\begin{aligned}
 I_1 &= \frac{(q_2 - 1)^2}{q_1^3 q_2^2 (q_2 - 1)} \int_{q_1^{-1} q_2^{-\frac{3}{2}} a\mathbb{T}} \frac{f(q_2^{\frac{3}{2}} s, q_2^{-1})}{c_2(s)c_2(s^{-1})} ds, \\
 I_2 &= \frac{(q_1 - 1)^2}{q_1 q_2^3 (q_1^2 - 1)} \int_{q_1^{-\frac{1}{2}} c\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}} s)}{c_1(s)c_1(s^{-1})} ds,
 \end{aligned}$$

$$\begin{aligned}
 I_3 &= \frac{(q_1 - 1)^2}{q_1 q_2^3 (q_1^2 - 1)} \int_{q_1^{-\frac{1}{2}} c \mathbb{T}} \frac{f(q_1^{\frac{1}{2}} s^{-3}, q_1^{-\frac{1}{2}} s)}{c_1(s) c_1(s^{-1})} ds, \\
 I_4 &= \frac{(q_2 - 1)^2}{q_1^3 q_2^2 (q_2^2 - 1)} \int_{q_2^{\frac{1}{2}} c \mathbb{T}} \frac{f(s^{-2}, q_2^{-\frac{1}{2}} s)}{c_2(s) c_2(s^{-1})} ds, \\
 I_5 &= \frac{(q_2 - 1)^2}{q_1^3 q_2^2 (q_2^2 - 1)} \int_{q_2^{-\frac{1}{2}} c \mathbb{T}} \frac{f(q_2^{-\frac{3}{2}} s^{-1}, q_2^{\frac{1}{2}} s)}{c_2(s) c_2(s^{-1})} ds, \\
 I_6 &= \frac{(q_1 - 1)^2}{q_1 q_2^3 (q_1^2 - 1)} \int_{c^{\frac{1}{2}} \mathbb{T}} \frac{f(q_1^{-\frac{1}{2}} s^{-3}, s^2)}{c_1(s) c_1(s^{-1})} ds,
 \end{aligned}$$

where we have put $s = q_2^{-\frac{3}{2}} t_1, q_1^{-\frac{1}{2}} t_2, q_1^{\frac{1}{2}} t_2, q_2^{\frac{1}{2}} t_2, q_2^{-\frac{1}{2}} t_2$, and t_2 in I_1, I_2, I_3, I_4, I_5 , and I_6 , respectively.

One now shifts each contour to \mathbb{T} . As we discuss below, some complications arise when $q_1 = q_2$ or $q_1 = q_2^3$, and so suppose for now that $q_1 \neq q_2$ and $q_1 \neq q_2^3$. As in the $\tilde{C}_2, L = Q$, case the integrands of I_1, \dots, I_6 are all nonsingular on \mathbb{T} . Moreover, assuming that $q_1 \neq q_2$ and $q_1 \neq q_2^3$, all singularities are simple poles, and at the special values $q_1^2 = q_2^3$ or $q_1 = q_2^2$ there are some removable singularities. Write I_1^u, \dots, I_6^u for the integrals over the contour \mathbb{T} . A lengthy analysis (using the fact that a is close to 0 and c is close to 1) gives

$$\begin{aligned}
 \text{Tr}(h) &= \frac{1}{q_1^3 q_2^3} \iint_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt + I_1^u + \dots + I_6^u + Af(q_1^{-1}, q_2^{-1}) + B_+ \sigma_+ + B_- \sigma_- \\
 &\quad + C(f(q_1^{-1}, \omega) + f(q_1^{-1}, \omega^{-1})) \\
 &\quad + |D| \times \begin{cases} f(q_1, q_2^{-1}) & \text{if } q_1 < q_2^{3/2}, \\ \sigma & \text{if } q_2^{3/2} < q_1 < q_2^2, \\ f(q_1^{-1}, q_2) & \text{if } q_2^2 < q_1, \end{cases}
 \end{aligned}$$

where $\sigma_{\pm} = f(\pm q_1^{-1/2} q_2^{3/2}, q_2^{-1}) + f(q_1^{-1}, \pm q_1^{1/2} q_2^{-1/2}) + f(\pm q_1^{1/2} q_2^{-3/2}, \pm q_1^{-1/2} q_2^{1/2})$ and

$$\begin{aligned}
 \sigma &= f(q_1^{-1} q_2^3, q_2^{-1}) + f(q_1^{-1}, q_1 q_2^{-1}) + f(q_1^2 q_2^{-3}, q_1^{-1} q_2) + f(q_1^{-2} q_2^3, q_1 q_2^{-2}) \\
 &\quad + f(q_1 q_2^{-3}, q_1^{-1} q_2^2).
 \end{aligned}$$

As in the previous sections, it is easy to show that

$$I_1^u + \dots + I_6^u = \frac{(q_1 - 1)^2}{2q_1 q_2^3 (q_1^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds + \frac{(q_2 - 1)^2}{2q_1^3 q_2^2 (q_2^2 - 1)} \int_{\mathbb{T}} \frac{\chi_s^2(h)}{|c_2(s)|^2} ds.$$

Proposition 3.10 gives $f(q_1^{-1}, \omega) + f(q_1^{-1}, \omega^{-1}) = \pi^8(h)$ and $\sigma = \pi^6(h)$ (note that π^6 only occurs for parameters $q_2^{3/2} < q_1 < q_2^2$, and in this range π^6 is defined and has regular central character). We also have $\sigma_{\pm} = \pi_{\pm}^7(h)$. For π_+^7 it is important that $q_1 \neq q_2$ and $q_1 \neq q_2^3$, for otherwise π_+^7 does not have a regular central character and things become complicated (see below). Lemma 3.11 gives $f(q_1^{-1}, q_2^{-1}) = \chi^3(h)$. Since we exclude $q_1 = q_2$, the representation π^4 has regular central character for parameters $q_1 < q_2^{3/2}$, and so Lemma 3.11 gives $f(q_1, q_2^{-1}) = \chi^4(h)$. Similarly, $f(q_1^{-1}, q_2) = \chi^5(h)$ for all $q_2^2 < q_1$ with $q_1 \neq q_2^3$.

It remains to discuss the cases $q_1 = q_2$ and $q_1 = q_2^3$. Let us briefly outline the work involved. Consider the $q_1 = q_2$ case (the $q_1 = q_2^3$ case is similar). The integrands of I_1, \dots, I_6 are still nonsingular on \mathbb{T} , and the contours in the integrals I_3, I_4 and I_6 can all be shifted to \mathbb{T} without encountering any poles. This leaves I_1, I_2 and I_5 to consider. Writing $f(t) = g(t)/d(t)$ the integrands of I_1, I_2 and I_5 are respectively

$$\frac{qs^2(1-s^2)g(q^{\frac{3}{2}}s, q^{-1})}{(1-q)(1-q^{-\frac{1}{2}}s^{-1})^2(1+q^{-\frac{1}{2}}s^{-1})(1-q^{-\frac{5}{2}}s^{-1})(1-q^{-\frac{1}{2}}s)^2(1+q^{-\frac{1}{2}}s)(1-q^{-\frac{5}{2}}s)},$$

$$\frac{s^4(1-s^2)g(q^{-1}, q^{\frac{1}{2}}s)}{(1-q)(1-q^{-\frac{3}{2}}s^{-3})(1-q^{-1}s^{-2})(1-q^{-\frac{3}{2}}s^{-1})(1-q^{-\frac{3}{2}}s^3)(1-q^{-1}s^2)(1-q^{-\frac{3}{2}}s)},$$

$$\frac{q^{-\frac{1}{2}}s(1-s^2)g(q^{-\frac{3}{2}}s^{-1}, q^{\frac{1}{2}}s)}{(q-1)(1-q^{-\frac{1}{2}}s^{-1})^2(1+q^{-\frac{1}{2}}s^{-1})(1-q^{-\frac{5}{2}}s^{-1})(1-q^{-\frac{1}{2}}s)^2(1+q^{-\frac{1}{2}}s)(1-q^{-\frac{5}{2}}s)}.$$

The relevant poles are at $s = q^{-\frac{1}{2}}$ (a double pole for I_1, I_2 , and I_5), $s = q^{-\frac{1}{2}}$ (a single pole for I_1, I_2 , and I_5), $s = \omega^{\pm 1}q^{-\frac{1}{2}}$ (single poles for I_2 only), and $s = q^{-\frac{5}{2}}$ (a single pole for I_1 only). The residue contributions from $s = -q^{-\frac{1}{2}}$ make up the $\chi_-^7(h)$ term, the contributions from $s = \omega^{\pm 1}q^{-\frac{1}{2}}$ give the $\chi^8(h)$ term, and the contribution from $s = q^{-\frac{5}{2}}$ gives the $\chi^3(h)$ term. All that remains is to analyse the contribution from the double poles of each integral at $s = q^{-\frac{1}{2}}$.

We claim that the combined residue contribution from the point $s = q^{-\frac{1}{2}}$ is

$$R_1 + R_2 + R_5 = \frac{q(q-1)^3}{6(q+1)^2(q^3-1)} (\chi_+^7(h) + 2\chi^4(h)). \tag{4.6}$$

We do not have a conceptual proof of this fact, but it can be obtained by direct calculation as follows. As in Remark 2.2, the functions $g(t) = g_t(h)$ can be explicitly computed (since one only needs to know the values $g_t(T_w)$ for $w \in W_0$, as $g_t(T_w x^\lambda) = t^\lambda g_t(T_w)$). Then the residue contributions can be explicitly calculated (making 12 separate calculations, one for each $h = T_w x^\lambda$ with $w \in W_0$). On the other hand, using the explicit matrices (Example 2 of Sect. 3.3) for the representations π_+^7 and π^4 one can compute the expression $\chi_+^7(T_w x^\lambda) + 2\chi^4(T_w x^\lambda)$ and compare. This completes the proof. □

4.7 The $\tilde{BC}_2(q_0, q_1, q_2)$ algebras with $L = Q$

The root system is $R = \pm\{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, \alpha_1 + 2\alpha_2, 2\alpha_2, 2(\alpha_1 + \alpha_2)\}$, giving dual root system $R^\vee = \pm\{\alpha_1^\vee, \alpha_2^\vee, \alpha_1^\vee + \alpha_2^\vee, 2\alpha_1^\vee + \alpha_2^\vee, \alpha_2^\vee/2, \alpha_1^\vee + \alpha_2^\vee/2\}$. The affine Hecke algebra has generators $T_1, T_2, x_1 = x^{\alpha_1^\vee}$ and $x_2 = x^{\alpha_2^\vee/2}$ with relations

$$\begin{aligned} T_1^2 &= 1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})T_1, & T_1x_1 &= x_1^{-1}T_1 + (q_1^{\frac{1}{2}} - q_1^{-\frac{1}{2}})(1 + x_1), \\ T_2^2 &= 1 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})T_2, & T_2x_2 &= x_2^{-1}T_2 + (q_2^{\frac{1}{2}} - q_2^{-\frac{1}{2}})x_2 + (q_0^{\frac{1}{2}} - q_0^{-\frac{1}{2}}), \\ T_1T_2T_1T_2 &= T_2T_1T_2T_1, & T_2x_1 &= x_1x_2^2T_2^{-1} - (q_0^{\frac{1}{2}} - q_0^{-\frac{1}{2}})x_1x_2, \\ x_1x_2 &= x_2x_1, & T_1x_2 &= x_1x_2T_1^{-1}. \end{aligned}$$

Let $\pi_t = \text{Ind}_{\mathbb{C}[Q]}^{\mathcal{H}}(\mathbb{C}v_t)$ be the principal series representation of \mathcal{H} with central character $t = (t_1, t_2) \in (\mathbb{C}^\times)^2$, where $\mathbb{C}v_t$ is the 1-dimensional representation of $\mathbb{C}[Q]$ with $x_1 \cdot v_t = t_1 v_t$ and $x_2 \cdot v_t = t_2 v_t$.

Let \mathcal{H}_1 be the subalgebra generated by T_1, x_1, x_2 and let \mathcal{H}_2 be the subalgebra generated by T_2, x_1, x_2 . Let $s \in \mathbb{C}^\times$, and let $\pi_s^1 = \text{Ind}_{\mathcal{H}_1}^{\mathcal{H}}(\mathbb{C}u_s^1)$ and $\pi_s^j = \text{Ind}_{\mathcal{H}_2}^{\mathcal{H}}(\mathbb{C}u_s^j)$ ($j = 2, 3, 4$) be the 4-dimensional representations induced from the 1-dimensional representation $\mathbb{C}u_s^1$ of \mathcal{H}_1 and the 1-dimensional representations $\mathbb{C}u_s^j$ ($j = 2, 3, 4$) of \mathcal{H}_2 given by

$$\begin{aligned} T_1 \cdot u_s^1 &= -q_1^{-\frac{1}{2}}u_s^1, & x_1 \cdot u_s^1 &= q_1^{-1}u_s^1, & x_2 \cdot u_s^1 &= q_1^{\frac{1}{2}}su_s^1, \\ T_2 \cdot u_s^2 &= -q_2^{-\frac{1}{2}}u_s^2, & x_1 \cdot u_s^2 &= q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}su_s^2, & x_2 \cdot u_s^2 &= q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}u_s^2, \\ T_2 \cdot u_s^3 &= -q_2^{-\frac{1}{2}}u_s^3, & x_1 \cdot u_s^3 &= q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}su_s^3, & x_2 \cdot u_s^3 &= -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}u_s^3, \\ T_2 \cdot u_s^4 &= q_2^{\frac{1}{2}}u_s^4, & x_1 \cdot u_s^4 &= q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}su_s^4, & x_2 \cdot u_s^4 &= -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}u_s^4. \end{aligned}$$

Let π^j ($j = 5, \dots, 11$) be the 1-dimensional representations of \mathcal{H} with

$$\begin{aligned} \pi^5 &= (-q_1^{-\frac{1}{2}}, -q_2^{-\frac{1}{2}}, q_1^{-1}, q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}), & \pi^6 &= (-q_1^{-\frac{1}{2}}, -q_2^{-\frac{1}{2}}, q_1^{-1}, -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}), \\ \pi^7 &= (q_1^{\frac{1}{2}}, -q_2^{-\frac{1}{2}}, q_1, q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}), & \pi^8 &= (q_1^{\frac{1}{2}}, -q_2^{-\frac{1}{2}}, q_1, -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}), \\ \pi^9 &= (-q_1^{-\frac{1}{2}}, q_2^{\frac{1}{2}}, q_1^{-1}, q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}), & \pi^{10} &= (-q_1^{-\frac{1}{2}}, q_2^{\frac{1}{2}}, q_1^{-1}, -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}), \\ \pi^{11} &= (q_1^{\frac{1}{2}}, q_2^{\frac{1}{2}}, q_1, -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}), \end{aligned}$$

where in each case we list the quadruples $(\pi^j(T_1), \pi^j(T_2), \pi^j(x_1), \pi^j(x_2))$.

Let $\pi^{12} = M_J(s), \pi^{13} = M_J(t)$, and $\pi^{14} = M_J(u)$ be the 3-dimensional representations with

$$\begin{aligned} (s^{\alpha_1^\vee}, s^{\alpha_2^\vee/2}) &= (q_1^{-1}, q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}), & (t^{\alpha_1^\vee}, t^{\alpha_2^\vee/2}) &= (q_1^{-1}, -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}), \\ (u^{\alpha_1^\vee}, u^{\alpha_2^\vee/2}) &= (q_1^{-1}, -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}), \end{aligned}$$

and $J = \{\alpha_2\}$. We assume that $q_1 \neq q_0q_2$ and $q_1^2 \neq q_0q_2$ for π^{12} , that $q_1 \neq q_0^{-1}q_2$ and $q_1^2 \neq q_0^{-1}q_2$ for π^{13} , and that $q_1 \neq q_0q_2^{-1}$ and $q_1^2 \neq q_0q_2^{-1}$ for π^{14} , so that $N(s) = N(t) = N(u) = \{\alpha_1, \alpha_2\}$ and $D(s) = D(t) = D(u) = \emptyset$, and hence $F_J(s) = F_J(t) = F_J(u) = \{s_2, s_1s_2, s_2s_1s_2\}$.

Finally, let $\pi^{15} = M_J(t)$ and $\pi^{16} = M_J(u)$ be the 2-dimensional representations with

$$(t^{\alpha_1^\vee}, t^{\alpha_2^\vee/2}) = (-q_0, q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}), \quad (u^{\alpha_1^\vee}, u^{\alpha_2^\vee/2}) = (-q_2, q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}), \quad J = \emptyset.$$

Theorem 4.8 *For all $h \in \mathcal{H}$ we have*

$$\begin{aligned} \text{Tr}(h) &= \frac{1}{8q_1^2q_2^2} \iint_{\mathbb{T}^2} \frac{\chi_t(h)}{|c(t)|^2} dt + C_6 \int_{\mathbb{T}} \frac{\chi_s^1(h)}{|c_1(s)|^2} ds + C_7 \int_{\mathbb{T}} \frac{\chi_s^2(h)}{|c_2(s)|^2} ds \\ &+ C_8 \int_{\mathbb{T}} \frac{\chi(h)}{|c_3(s)|^2} ds + C_1 \chi^5(h) \\ &+ |C_2| \times \begin{cases} \pi^7(h) & \text{if } q_1 < q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}, \\ \pi^{12}(h) & \text{if } q_0^{\frac{1}{2}}q_2^{\frac{1}{2}} < q_1 < q_0q_2, \\ \pi^9(h) & \text{if } q_0q_2 < q_1 \end{cases} + \begin{cases} X_1 & \text{if } q_0 < q_2, \\ X_2 & \text{if } q_2 < q_0, \end{cases} \end{aligned}$$

where $\chi(h) = \chi_s^3(h)$ if $q_0 < q_2$ and $\chi(h) = \chi_s^4(h)$ if $q_2 < q_0$, and where

$$\begin{aligned} X_1 &= |C_3| \chi^{15}(h) + |C_4| \chi^6(h) + |C_5| \times \begin{cases} \chi^8(h) & \text{if } q_1 < q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}, \\ \chi^{13}(h) & \text{if } q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}} < q_1 < q_0^{-1}q_2, \\ \pi^{10}(h) & \text{if } q_0^{-1}q_2 < q_1, \end{cases} \\ X_2 &= |C_3| \chi^{16}(h) + |C_5| \pi^{10}(h) + |C_4| \times \begin{cases} \pi^{11}(h) & \text{if } q_1 < q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}, \\ \pi^{14}(h) & \text{if } q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}} < q_1 < q_0q_2^{-1}, \\ \pi^6(h) & \text{if } q_0q_2^{-1} < q_1, \end{cases} \end{aligned}$$

with $c(t), c_1(s), c_2(s), c_3(s), C_1, \dots, C_8$ as in Appendix A.4.

Proof The series $G_t(h)$ converges for $|t_1| < q_1^{-1}$ and $|t_2| < q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}$, and so writing $f(t) = f_t(h)$ we have

$$\text{Tr}(h) = \frac{1}{q_1^2q_2^2} \int_{q_1^{-1}a\mathbb{T}} \int_{q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}b\mathbb{T}} \frac{f(t)}{c(t)c(t^{-1})} dt_2 dt_1$$

whenever $0 < a, b < 1$. We choose a and b both very close to 0, and choose $0 < c < 1$ very close to 1. The t_2 -poles of the integrand between the contour $q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}b\mathbb{T}$ and

$c\mathbb{T}$ are at $t_2 = q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}$, at $t_2 = -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}$ (if $q_0 < q_2$) and at $t_2 = -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}$ (if $q_2 < q_0$). Thus we can shift the t_2 -contour to $c\mathbb{T}$ at the cost of residue contributions from the above points. Now interchange the order of integration in the double integral. The t_1 -poles of the integrand between $q_1^{-1}a\mathbb{T}$ and \mathbb{T} are at $t_1 = q_1^{-1}$, $t_1 = q_1^{-1}t_2^{-2}$, $t_1 = q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}t_2^{-1}$, $t_1 = -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}t_2^{-1}$ (if $q_0 < q_2$) and $t_1 = -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}t_2^{-1}$ (if $q_2 < q_0$). Computing the associated residues gives

$$\text{Tr}(h) = \frac{1}{q_1^2 q_2^2} \int_{\mathbb{T}^2} \frac{f(t)}{|c(t)|^2} dt + I_1 + I_2 + I_3 + I_4 + \begin{cases} I_5 + I_6 & \text{if } q_0 < q_2, \\ I'_5 + I'_6 & \text{if } q_2 < q_0, \end{cases}$$

where

$$I_1 = \frac{q_0 q_2 - 1}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{q_0^{-\frac{1}{2}}q_1^{-1}q_2^{-\frac{1}{2}}a\mathbb{T}} \frac{f(q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}s, q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}})}{c_2(s)c_2(s^{-1})} ds,$$

$$\text{where } s = q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}t_1,$$

$$I_2 = \frac{q_1 - 1}{q_1 q_2^2 (q_1 + 1)} \int_{q_1^{-\frac{1}{2}}c\mathbb{T}} \frac{f(q_1^{-1}, q_1^{\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds, \quad \text{where } s = q_1^{-\frac{1}{2}}t_2,$$

$$I_3 = \frac{q_1 - 1}{q_1 q_2^2 (q_1 + 1)} \int_{q_1^{\frac{1}{2}}c\mathbb{T}} \frac{f(s^{-2}, q_1^{-\frac{1}{2}}s)}{c_1(s)c_1(s^{-1})} ds, \quad \text{where } s = q_1^{\frac{1}{2}}t_2,$$

$$I_4 = \frac{q_0 q_2 - 1}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{c\mathbb{T}} \frac{f(q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1}, s)}{c_2(s)c_2(s^{-1})} ds, \quad \text{where } s = t_2,$$

$$I_5 = \frac{q_2 - q_0}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{q_0^{\frac{1}{2}}q_1^{-1}q_2^{-\frac{1}{2}}a\mathbb{T}} \frac{f(q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}s, -q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}})}{c_3(s)c_3(s^{-1})} ds,$$

$$\text{where } s = q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}t_1,$$

$$I'_5 = \frac{q_0 - q_2}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{q_0^{-\frac{1}{2}}q_1^{-1}q_2^{\frac{1}{2}}a\mathbb{T}} \frac{f(q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}s, -q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}})}{c_3(s)c_3(s^{-1})} ds,$$

$$\text{where } s = q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}t_1,$$

$$I_6 = \frac{q_2 - q_0}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{c\mathbb{T}} \frac{f(q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1}, -s)}{c_3(s)c_3(s^{-1})} ds, \quad \text{where } s = -t_2,$$

$$I'_6 = \frac{q_0 - q_2}{q_1^2 q_2^2 (q_0 + 1)(q_2 + 1)} \int_{c\mathbb{T}} \frac{f(q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}s^{-1}, -s)}{c_3(s)c_3(s^{-1})} ds, \quad \text{where } s = -t_2.$$

Now shift the contours in all integrals I_j, I'_j to \mathbb{T} . We omit the details of this long calculation. □

Acknowledgements Partially supported under the Australian Research Council (ARC) discovery grant DP110103205.

Appendix: Constants and c-functions

We write $\sigma_1(x) = 1 + x$ and $\sigma_2(x) = 1 + x + x^2$.

A.1 $\tilde{C}_2(q_1, q_2)$ algebras with $L = Q$

$$c(t) = \frac{(1 - q_1^{-1}t_1^{-1})(1 - q_1^{-1}t_1^{-1}t_2^{-2})(1 - q_2^{-1}t_2^{-1})(1 - q_2^{-1}t_1^{-1}t_2^{-1})}{(1 - t_1^{-1})(1 - t_1^{-1}t_2^{-2})(1 - t_2^{-1})(1 - t_1^{-1}t_2^{-1})},$$

$$c_1(s) = \frac{(1 + q_1^{-\frac{1}{2}}s^{-1})(1 - q_1^{-\frac{1}{2}}q_2^{-1}s^{-1})(1 - q_1^{\frac{1}{2}}q_2^{-1}s^{-1})}{(1 - s^{-2})(1 - q_1^{\frac{1}{2}}s^{-1})},$$

$$c_2(s) = \frac{(1 - q_1^{-1}q_2^{-1}s^{-1})(1 - q_1^{-1}q_2s^{-1})}{(1 - s^{-1})(1 - q_2s^{-1})},$$

$$A = \frac{(q_1q_2 - 1)(q_1q_2^2 - 1)}{\sigma_1(q_1)\sigma_1(q_2)^2\sigma_1(q_1q_2)}, \quad B = \frac{2q_2(q_1 - 1)^2}{\sigma_1(q_2)^2\sigma_1(q_1q_2^{-1})\sigma_1(q_1q_2)},$$

$$C = \frac{(q_1q_2^{-1} - 1)(1 - q_1q_2^{-2})}{\sigma_1(q_1)\sigma_1(q_2^{-1})^2\sigma_1(q_1q_2^{-1})}.$$

A.2 $\tilde{C}_2(q_1, q_2)$ algebras with $L = P$

$$c(t) = \frac{(1 - q_1^{-1}t_1^{-2}t_2^2)(1 - q_1^{-1}t_2^{-2})(1 - q_2^{-1}t_1t_2^{-2})(1 - q_2^{-1}t_1^{-1})}{(1 - t_1^{-2}t_2^2)(1 - t_2^{-2})(1 - t_1t_2^{-2})(1 - t_1^{-1})},$$

$$c_1(s) = \frac{(1 + q_1^{-\frac{1}{2}}s^{-1})(1 - q_1^{-\frac{1}{2}}q_2^{-1}s^{-1})(1 - q_1^{\frac{1}{2}}q_2^{-1}s^{-1})}{(1 - s^{-2})(1 - q_1^{\frac{1}{2}}s^{-1})},$$

$$c_2(s) = \frac{(1 - q_1^{-1}q_2^{-1}s^{-2})(1 - q_1^{-1}q_2s^{-2})}{(1 - s^{-2})(1 - q_2s^{-2})}.$$

A.3 $\tilde{G}_2(q_1, q_2)$ algebras with $L = Q$

$$c(t) = \left((1 - q_1^{-1}t_1^{-1})(1 - q_1^{-1}t_1^{-2}t_2^{-3})(1 - q_1^{-1}t_1^{-1}t_2^{-3})(1 - q_2^{-1}t_2^{-1}) \right. \\ \times (1 - q_2^{-1}t_1^{-1}t_2^{-2})(1 - q_2^{-1}t_1^{-1}t_2^{-1}) \left. \right) \left((1 - t_1^{-1})(1 - t_1^{-2}t_2^{-3})(1 - t_1^{-1}t_2^{-3}) \right. \\ \left. \times (1 - t_2^{-1})(1 - t_1^{-1}t_2^{-2})(1 - t_1^{-1}t_2^{-1}) \right)^{-1},$$

$$c_1(s) = \frac{(1 - q_1^{-\frac{1}{2}}\omega s^{-1})(1 - q_1^{-\frac{1}{2}}\omega^{-1}s^{-1})(1 - q_2^{-1}s^{-2})(1 - q_1^{-\frac{1}{2}}q_2^{-1}s^{-1})(1 - q_1^{\frac{1}{2}}q_2^{-1}s^{-1})}{(1 - s^{-2})(1 - q_1^{-\frac{1}{2}}s^{-1})(1 - q_1^{\frac{1}{2}}s^{-3})},$$

$$c_2(s) = \frac{(1 - q_1^{-1}s^{-2})(1 - q_1^{-1}q_2^{-\frac{3}{2}}s^{-1})(1 - q_1^{-1}q_2^{\frac{3}{2}}s^{-1})}{(1 - s^{-2})(1 - q_2^{\frac{3}{2}}s^{-1})(1 - q_2^{\frac{1}{2}}s^{-1})},$$

$$A = \frac{(q_1q_2^2 - 1)(q_1^2q_2^3 - 1)}{\sigma_1(q_1)\sigma_1(q_2)\sigma_2(q_2)\sigma_2(q_1q_2)},$$

$$B_{\pm} = \frac{q_1(q_1 - 1)(q_2 - 1)}{2\sigma_1(q_1)\sigma_1(q_2)\sigma_2(\pm\sqrt{q_1/q_2})\sigma_2(\pm\sqrt{q_1q_2})},$$

$$C = \frac{q_2(q_1 - 1)(q_1^3 - 1)}{\sigma_2(q_2)\sigma_2(q_1q_2^{-1})\sigma_2(q_1q_2)}, \quad D = \frac{(1 - q_1q_2^{-2})(q_1^2q_2^{-3} - 1)}{\sigma_1(q_1)\sigma_1(q_2^{-1})\sigma_2(q_2^{-1})\sigma_2(q_1q_2^{-1})}.$$

A.4 $\tilde{BC}_2(q_0, q_1, q_2)$ algebras with $L = Q$

$$c(t) = \frac{(1 - q_1^{-1}t_1^{-1})(1 - q_1^{-1}t_1^{-1}t_2^{-2})(1 - a^{-1}t_1^{-1}t_2^{-1})(1 + b^{-1}t_1^{-1}t_2^{-1})(1 - a^{-1}t_2^{-1})(1 + b^{-1}t_2^{-1})}{(1 - t_1^{-1})(1 - t_1^{-1}t_2^{-2})(1 - t_1^{-2}t_2^{-2})(1 - t_2^{-2})},$$

$$c_1(s) = \frac{(1 - q_0^{-\frac{1}{2}}q_1^{-\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})(1 + q_0^{\frac{1}{2}}q_1^{-\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})(1 - q_0^{-\frac{1}{2}}q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})(1 + q_0^{\frac{1}{2}}q_1^{\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})}{(1 - s^{-2})(1 - q_1s^{-2})},$$

$$c_2(s) = \frac{(1 + q_0^{\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})(1 - q_0^{-\frac{1}{2}}q_1^{-1}q_2^{-\frac{1}{2}}s^{-1})(1 - q_0^{\frac{1}{2}}q_1^{-1}q_2^{\frac{1}{2}}s^{-1})}{(1 - s^{-2})(1 - q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}s^{-1})},$$

$$c_3(s) = \frac{(1 + q_0^{-\frac{1}{2}}q_2^{-\frac{1}{2}}s^{-1})(1 - q_0^{\frac{1}{2}}q_1^{-1}q_2^{-\frac{1}{2}}s^{-1})(1 - q_0^{-\frac{1}{2}}q_1^{-1}q_2^{\frac{1}{2}}s^{-1})}{(1 - s^{-2})(1 - q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}s^{-1})},$$

where $a = q_0^{\frac{1}{2}}q_2^{\frac{1}{2}}$ and $b = q_0^{-\frac{1}{2}}q_2^{\frac{1}{2}}$.

$$C_1 = \frac{(q_0q_1q_2 - 1)(q_0q_1^2q_2 - 1)}{\sigma_1(q_0)\sigma_1(q_1)\sigma_1(q_2)\sigma_1(q_0q_1)\sigma_1(q_1q_2)},$$

$$C_3 = \frac{(q_2 - q_0)(q_0q_2 - 1)}{\sigma_1(q_0q_1^{-1})\sigma_1(q_1^{-1}q_2)\sigma_1(q_0q_1)\sigma_1(q_1q_2)},$$

and $C_2 = -C_1(q_0^{-1}, q_1, q_2^{-1})$, $C_4 = C_1(q_0^{-1}, q_1, q_2)$ and $C_5 = -C_1(q_0, q_1, q_2^{-1})$. Finally,

$$C_6 = \frac{q_1 - 1}{2q_1q_2^2(q_1 + 1)}, \quad C_7 = \frac{q_0q_2 - 1}{2q_1^2q_2(q_0 + 1)(q_2 + 1)},$$

$$C_8 = \frac{|q_2 - q_0|}{2q_1^2q_2(q_0 + 1)(q_2 + 1)}.$$

References

1. Aubert, A.M., Plymen, R.: Plancherel measure for $GL(n, F)$ and $GL(m, D)$: explicit formulas and Bernstein decomposition. *J. Number Theory* **112**, 26–66 (2005)
2. Borel, A.: Admissible representations of a semisimple group over a local field with fixed vectors under an Iwahori subgroup. *Invent. Math.* **35**, 233–259 (1976)
3. Bourbaki, N.: *Lie Groups and Lie Algebras, Chapters 4–6*. Elements of Mathematics. Springer, Berlin (2002)
4. Ciubotaru, D., Kato (Shiota), M., Kato, S.: On characters and formal degrees of discrete series of affine Hecke algebras of classical types. *Invent. Math.* (2011). doi:[10.1007/s00222-011-0338-3](https://doi.org/10.1007/s00222-011-0338-3). Online
5. Davis, M.: Representations of rank 2 affine Hecke algebras at roots of unity (2011). [arXiv:1104.4826v1](https://arxiv.org/abs/1104.4826v1)
6. Delorme, P., Opdam, E.: The Schwartz algebra of an affine Hecke algebra. *J. Reine Angew. Math.* **625**, 59–114 (2008)
7. Dixmier, J.: *C^* -Algebras*. North-Holland Mathematical Library, vol. 15. North-Holland, Amsterdam (1977)
8. Geck, M., Pfeiffer, G.: *Characters of Finite Coxeter Groups and Iwahori–Hecke Algebras* London Mathematical Society Monographs, New Series, vol. 21. Clarendon Press, Oxford (2000)
9. Harish-Chandra: In: Varadarajan, V.S. (ed.) *Collected Papers, Vol. IV, 1970–1983*. Springer, New York (1984)
10. Heckman, G., Opdam, E.: Harmonic Analysis for Affine Hecke Algebras. *Current Developments in Mathematics*, pp. 37–60. International Press, Somerville (1997)
11. Kato, S.-I.: Irreducibility of principal series representations for Hecke algebras of affine type. *J. Fac. Sci., Univ. Tokyo, Sect. 1A, Math.* **28**(3), 929–943 (1981)
12. Kazhdan, D., Lusztig, G.: Proof of the Deligne–Langlands conjecture for Hecke algebras. *Invent. Math.* **87**, 153–215 (1987)
13. Kutzko, P., Morris, L.: Explicit Plancherel theorems for $H(q_1, q_2)$ and $SL_2(F)$. *Pure Appl. Math. Q.* **5**, 435–467 (2009)
14. Lusztig, G.: Affine Hecke algebras and their graded versions. *J. Am. Math. Soc.* **2**(3), 599–635 (1989)
15. Macdonald, I.G.: *Affine Hecke Algebras and Orthogonal Polynomials*. Cambridge Tracts in Mathematics, vol. 157. Cambridge University Press, Cambridge (2003)
16. Matsumoto, H.: *Analyse Harmonique dans les Systèmes de Tits Bornologiques de Type Affine*. Lecture Notes in Mathematics, vol. 590. Springer, Berlin (1977)
17. Nelson, K., Ram, A.: Kostka–Foulkes polynomials and Macdonald spherical functions. In: *Surveys in Combinatorics, 2003 (Bangor)*. London Math. Soc. Lecture Note Ser., vol. 307, pp. 325–370. Cambridge University Press, Cambridge (2003)
18. Opdam, E.: A generating function for the trace of the Iwahori–Hecke algebra. In: *Studies in Memory of Issai Schur (Chevaleret/Rehovot, 2000)*. *Progr. Math.*, vol. 210, pp. 301–323. Birkhäuser, Boston (2003)
19. Opdam, E.: On the spectral decomposition of affine Hecke algebras. *J. Inst. Math. Jussieu* **3**(4), 531–648 (2004)
20. Opdam, E.: The central support of the Plancherel measure of an affine Hecke algebra. *Mosc. Math. J.* **7**(4), 723–741, 767–768 (2007)
21. Opdam, E., Solleveld, M.: Discrete series characters for affine Hecke algebras and their formal degrees. *Acta Math.* **205**(1), 105–187 (2010)
22. Parkinson, J., Schapira, B.: A local limit theorem for random walk on the chambers of \tilde{A} buildings. In: *Random Walks, Boundaries and Spectra*. *Progress in Probability*, vol. 64, pp. 15–53 (2011)
23. Ram, A.: Calibrated representations of affine Hecke algebras (1998). [arXiv:math/0401323](https://arxiv.org/abs/math/0401323). Preprint, see also [24]
24. Ram, A.: Affine Hecke algebras and generalized standard Young tableaux. *J. Algebra* **230**, 367–415 (2003)
25. Ram, A.: Representations of rank two affine Hecke algebras. In: Musili, C. (ed.) *Advances in Algebra and Geometry*, pp. 57–91 (2003). Hindustan Book Agency
26. Reeder, M.: Hecke algebras and harmonic analysis on p -adic groups. *Am. J. Math.* **119**(1), 225–249 (1997)
27. Ronan, M.: *Lectures on Buildings, Revised Edn*. University of Chicago Press, Chicago (2009)
28. Steinberg, R.: Endomorphisms of Linear Algebraic Groups. *Mem. Am. Math. Soc.*, vol. 80. Amer. Math. Soc., Providence (1968)
29. Varadarajan, V.S.: *An Introduction to Harmonic Analysis on Semisimple Lie Groups*. Cambridge Studies in Advanced Mathematics, vol. 16. Cambridge University Press, Cambridge (1989)